arXiv:1203.0328v2 [math.AG] 17 May 2012

SCHUR FLEXIBILITY OF COMINUSCULE SCHUBERT VARIETIES

C. ROBLES

Abstract. Let $X = G/P$ be cominuscule rational homogeneous variety. (Equivalently, X admits the structure of a compact Hermitian symmetric space.) We say a Schubert class ξ is Schur rigid if the only irreducible subvarieties $Y \subset X$ with homology class $[Y] \in \mathbb{Z}\xi$ are Schubert varieties. Robles and The [18] identified a sufficient condition for ξ to be Schur rigid. In this paper we show that the condition is also necessary.

1. Introduction

Let $X = G/P$ be a rational homogeneous variety. The Schubert classes form an additive basis of the integral homology $H_\bullet(X, \mathbb{Z})$, [2]. In 1961, by Borel and Haefliger [3] asked: given a Schubert class ξ_w represented by a Schubert variety Y_w, aside from the G–translates $g \cdot Y_w$, are there any other algebraic representatives of the Schubert class? In some cases, the Schubert varieties are the only algebraic representatives, and we then say the Schubert class is *rigid*. This problem has been studied by several people, including R. Hartshorne, E. Rees and E. Thomas [9], R. Bryant [4], M. Walters [21], J. Hong [10, 11], and I. Coskun [6, 7].

It is a striking consequence of B. Kostant's work [14, 15] that, *when X is cominuscule*,* the varieties $Y \subset X$ that are homologous to an integer multiple of the Schubert class ξ_w (that is, $[Y] \in \mathbb{Z}\xi_w$) are characterized by a system of differential equations known as the *Schur system*, cf. Section 2.2. The Schubert varieties $\{g \cdot Y_w \mid g \in G\}$ are the *trivial solutions*. When there exist no nontrivial, irreducible solutions, we say that the Schubert class is *Schur rigid*. Every Schur rigid ξ_w is rigid, and this provides a differential-geometric approach to the algebro-topological question above.

Associated to the differential system is a Lie algebra cohomology. The cohomology contains a distinguished subspace $\mathcal{O}_w$. It is known [18] that the Schubert class ξ_w is Schur rigid when $\mathcal{O}_w = 0$; we say that the vanishing of $\mathcal{O}_w$ is a *cohomological obstruction to flexibility*. The goal of this paper is to show the converse: in the absence cohomological obstructions (equivalently, when $\mathcal{O}_w$ is nontrivial), there exist nontrivial solutions Y; precisely, Y is an irreducible subvariety of X such that (i) $[Y] \in \mathbb{Z}\xi_w$, and (ii) Y is not a G–translate of Y_w.

Date: February 24, 2013.

2010 *Mathematics Subject Classification.* 53C24, 53C30, 58A15, 58A17, 14M15, 14M17,

Key words and phrases. Rational homogeneous variety, cominuscule, compact Hermitian symmetric space, Schubert class, Schubert variety, Schur rigidity.

Robles is partially supported by NSF DMS-1006353.

*The Grassmannian $\mathrm{Gr}(k, n)$ of complex k–planes in $\mathbb{C}^n$ is an example of a cominuscule rational homogeneous variety (via the Plücker embedding).

1.1. **History.** Schur rigidity and the associated differential system were first studied independently by M. Walters [21] and R. Bryant [4]. Walters identified Schur rigid classes represented by (a) smooth Schubert varieties in $X = \mathrm{Gr}(k,n)$; and (b) codimension two Schubert varieties in $X = \mathrm{Gr}(2,n)$. Bryant identified Schur rigid classes represented by (a) smooth Schubert varieties in the cases that X is the Grassmannian $\mathrm{Gr}(k,n)$ or the Lagrangian Grassmannian $\mathrm{LG}(n,2n)$; (b) the maximal linear subspaces in the classical cominuscule X; and (c) singular Schubert varieties of low (co)dimension in $\mathrm{Gr}(k,n)$. In the case that X_w is smooth, the results of Bryant and Walters, which were obtained case-by-case, were generalized to arbitrary cominuscule X and given a beautiful, uniform proof by J. Hong [11]. There, the obstructions to flexibility are realized as a (Lie algebra) cohomological condition. Hong also showed that a large class of the singular Schubert varieties in the Grassmannian are Schur rigid [10]. C. Robles and D. The [18] extended the approach of [11] to give a complete list of the Schubert classes for which there exist cohomological obstructions to flexibility.

1.2. **Contents.** The main result of this paper is Theorem 3.1, which implies (Corollary 3.2) that in the absence of cohomological obstructions, the Schubert class is Schur flexible. As a consequence, we find that a Schubert class is Schur rigid if and only if its Poincaré dual is Schur rigid (Corollary 3.3).

Theorem 3.1 is proved by constructing nontrivial, irreducible integral varieties Y of the Schur system. The construction is guided by representation theoretic data associated to the Lie algebra cohomology group. While the representation theoretic construction is explicit, it is not clear, geometrically, what the integrals are. So, in Section 4, I give algebro-geometric descriptions of the integrals (Propositions 4.2, 4.5, 4.9, 4.13, 4.20). As an integral of the Schur system the homology class of Y satisfies $[Y] = r\xi_w$ for some integer $0 < r = r(Y)$. The descriptions of Section 4 allow us to determine the integers $r(Y)$.

Contents

1. Introduction 1
2. Review 3
Figure 1: Hasse poset and (a,J) data for the Cayley plane E_6/P_6 13
Figure 2: Hasse poset and (a,J) data for the Freudenthal variety E_7/E_7 14
3. Flexibility 15
4. The Schubert integrals 26
5. The Schur system 41
References 43

1.3. **Use of LiE.** There are two exceptional, irreducible compact Hermitian symmetric spaces: the Cayley plane E_6/P_6 and the Freudenthal variety E_7/P_7. The software [16] is used to perform computations and verify some results for these two cominuscule varieties. (Sections 2.12, 4.5 and 5.5.) The code I wrote applies to any cominuscule variety; that is, it is *not* specialized to the exceptional cases. This allowed me to test the code by applying it to classical cominuscule varieties (of low rank – the code requires that the rank of G be specified). The outputs are consistent with the "by hand" results obtained for the classical cases. This provides some confidence for the accuracy of the code and the corresponding claims in the exceptional cases.

1.4. **Acknowledgements.** Over the course of this project, I have benefitted from conversations and/or correspondence with many people, including R. Bryant, I. Coskun, J.M. Landsberg, N. Ressayre, F. Sottile and D. The. I thank them for their insights and time.

2. Review

2.1. Notation and background.

This is a continuation of [18]. With the exception noted in Remark 2.16, I will use the notation of that paper. To streamline the presentation, I will assume that the reader has reviewed the discussion of rational homogeneous varieties, their Schubert subvarieties, grading elements and Hasse diagrams in [18, Sections 2.1-2.4 and 3.1]. Briefly, G is a complex simple Lie group. A choice of Cartan and Borel subgroups $H \subset B$ has been fixed, $P \supset B$ is a maximal parabolic subgroup associated with a cominuscule root, and $X = G/P$ is the corresponding cominuscule variety. The associated Lie algebras are denoted $\mathfrak{h} \subset \mathfrak{b} \subset \mathfrak{p} \subset \mathfrak{g}$. Let W denote the Weyl group of $\mathfrak{g}$, and $W_\mathfrak{p}$ the Weyl group of the reductive component in the Levi decomposition of $\mathfrak{p}$. The Hasse diagram $W^\mathfrak{p}$ is the set of minimal length representatives of the coset space $W_\mathfrak{p}\backslash W$, and indexes the Schubert classes. Let

$$o = P/P \ \in \ X = G/P\,.$$

Given $w \in W^\mathfrak{p}$, the Zariski closure

$$Y_w \ := \ \overline{Bw^{-1}\cdot o}$$

is a Schubert variety. Any G–translate of the Schubert variety Y_w will be referred to as a *Schubert variety of type* w. Let $\xi_w = [Y_w] \in H_{2|w|}(X,\mathbb{Z})$ denote the corresponding Schubert class.

Let $\{Z_1,\ldots,Z_r\}$ be the basis of $\mathfrak{h}$ dual to the simple roots $\{\alpha_1,\ldots,\alpha_r\}$. Let $\alpha_\mathtt{i}$ be the simple root associated with the cominuscule $\mathfrak{p}$. Then

$$\mathfrak{g} \ = \ \mathfrak{g}_1 \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_{-1}\,, \quad \text{where} \quad \mathfrak{g}_k \ := \ \{A \in \mathfrak{g} \mid [Z_\mathtt{i}\,,\, A] = kA\}\,, \tag{2.1}$$

is the $Z_\mathtt{i}$*–graded decomposition* of the Lie algebra $\mathfrak{g}$. Moreover,

$$\mathfrak{p} \ = \ \mathfrak{g}_1 \oplus \mathfrak{g}_0\,, \tag{2.2}$$

and $\mathfrak{g}_0$ is the reductive component of the parabolic subalgebra $\mathfrak{p}$.

Remark 2.3. As a graded decomposition, we have $[\mathfrak{g}_k\,,\,\mathfrak{g}_\ell] \subset \mathfrak{g}_{k+\ell}$. In particular, the subalgebras $\mathfrak{g}_{\pm1} \subset \mathfrak{g}$ are abelian, $[\mathfrak{g}_1\,,\,\mathfrak{g}_1] \ = \ \{0\} \ = \ [\mathfrak{g}_{-1}\,,\,\mathfrak{g}_{-1}]$.

Let Δ denote the set of roots of $\mathfrak{g}$. Given $\alpha \in \Delta$, let $\mathfrak{g}_\alpha \subset \mathfrak{g}$ denote the corresponding root space. Given any subset $\mathfrak{s} \subset \mathfrak{g}$, let

$$\Delta(\mathfrak{s}) \ = \ \{\alpha \in \Delta \mid \mathfrak{g}_\alpha \subset \mathfrak{s}\}\,.$$

Given a subset U of a vector space, let $\langle U\rangle$ denote the linear span.

2.2. **The Schur differential system.** There is a natural $\mathfrak{g}_0$–module identification

$$T_oX \ \simeq \ \mathfrak{g}_{-1}\,.$$

The ℓ–th exterior power decomposes as

$$\textstyle\bigwedge^\ell T_oX \ = \ \bigoplus_{\substack{w\in W^{\mathfrak{p}}\\ |w|=\ell}} \mathbf{I}_w$$

into irreducible $\mathfrak{g}_0$–modules. (Cf. [18, Section 4.1].) The highest weight line $\mathfrak{n}_w \in \mathbb{P}\mathbf{I}_w$ is defined by

$$\Delta(w) \ = \ w\Delta^- \cap \Delta^+ \quad \text{and} \quad \mathfrak{n}_w \ = \ \bigoplus_{\alpha\in\Delta(w)} \mathfrak{g}_{-\alpha} \tag{2.4}$$

Remark 2.5. One important consequence of Remark 2.3 is that given a set $\Phi \subset \Delta(\mathfrak{g}_1)$, there exists $w \in W^{\mathfrak{p}}$ such that $\Delta(w) = \Phi$ if and only if $\Delta^+\backslash\Phi$ is closed. For details see [18, Section 2.3].

Consider the set of tangent ℓ–planes $\mathrm{Gr}(\ell, T_oX)$ as a subvariety of $\mathbb{P}(\bigwedge^\ell T_oX)$ via the Plücker embedding, and set $R_w = \mathrm{Gr}(|w|, T_oX) \cap \mathbb{P}\,\mathbf{I}_w$. The *Schur system* is the homogeneous bundle $\mathcal{R}_w \subset \mathrm{Gr}(|w|, TX)$ with fibre R_w over $o \in X$. An *integral manifold M of the Schur system* is a complex submanifold $M \subset X$ with the property that $TM \subset \mathcal{R}_w$. The *Schur system is rigid* if every connected integral manifold is contained in a Schubert variety of type w. An integral manifold that is not contained in any Schubert variety of type w we call *nontrivial.* When there exist nontrivial integrals we say the *Schur system is flexible.*

An *integral variety of the Schur system* is a subvariety $Y \subset X$ with the property that the set of smooth points Y^0 is an integral manifold of the Schur system. The Schubert class ξ_w (or the Schubert variety Y_w) is *Schur rigid* if every irreducible integral variety of the Schur system is a Schubert variety of type w. An irreducible integral variety that is not Schubert variety of type w is *nontrivial.* When there exist nontrivial integral varieties we say the Schubert class ξ_w (or the Schubert variety Y_w) is *Schur flexible.*

Theorem 2.6 ([4, 21])**.** *The homology class $[Y] \subset H_{2|w|}(X,\mathbb{Z})$ represented by a variety $Y \subset X$ is an integer multiple of the Schubert class ξ_w if and only if Y is an integral of the Schur system $\mathcal{R}_w$.*

Remark 2.7. *A priori*, it may happen that $\mathcal{R}_w$ admits nontrivial integral manifolds (a differential-geometric property), but no nontrivial integral varieties (an algebraic-geometric property). That is, it may be the case that $\mathcal{R}_w$ is flexible, while ξ_w is Schur rigid. However, we will see *a posteriori* (Corollary 3.2) that $\mathcal{R}_w$ is flexible if and only if ξ_w is Schur flexible.

For more on the Schur system see [18, Section 7] and the references therein.

2.3. **The Schubert differential system.** Recall the $Z_{\mathtt{i}}$–graded decomposition (2.1) of $\mathfrak{g}$, and let

$$G_0 \ := \ \{g \in G \mid \mathrm{Ad}_g(\mathfrak{g}_j) \subset \mathfrak{g}_j\}\,. \tag{2.8}$$

Then G_0 is a closed subgroup of G with Lie algebra $\mathfrak{g}_0$. Associated to the Schur system is the more restrictive *Schubert system* $\mathcal{B}_w$ defined by the G_0–orbit $B_w \subset \mathrm{Gr}(|w|, T_oX)$ of the $\mathfrak{g}_0$–highest weight line $\mathfrak{n}_w \in R_w$. In particular, $B_w \subset R_w$. The corresponding homogeneous bundle $\mathcal{B}_w \subset \mathrm{Gr}(|w|, TX)$ is precisely the set of $|w|$–planes tangent to (a smooth point of) a Schubert variety of type w. The notions of integrals, rigidity and flexibility for the Schubert system are analogous to those for the Schur system (Section 2.2). Theorem 2.6 and $\mathcal{B}_w \subset \mathcal{R}_w$ yield

Corollary 2.9. *The homology class $[Y] \subset H_{2|w|}(X, \mathbb{Z})$ represented by an integral variety $Y \subset X$ of the Schubert system $\mathcal{B}_w$ is an integer multiple of the Schubert class ξ_w.*

For more on the Schubert system see [18, Section 4] and the references therein. The following is [18, Proposition 7.5], and is due to Bryant (in the case that X is a Grassmannian) and Hong (in general).

Proposition 2.10. *The Schur system $\mathcal{R}_w$ is rigid if and only if $B_w = R_w$ and the Schubert system $\mathcal{B}_w$ is rigid.*

Remark 2.11. *(a) A posteriori* the condition that $R_w = B_w$ may be dropped; it is a consequence of Theorem 3.1 and Corollary 3.2 that $\mathcal{B}_w$ is rigid if and only if $\mathcal{R}_w$ is rigid.

(b) Note that R_w is the intersection of $\mathrm{Gr}(|w|, T_oX)$ with the (projective) linear span of B_w in $\mathbb{P}(\bigwedge^{|w|} T_oX)$. In general the containment $B_w \subset R_w$ is strict. In [18, Sections 8.2–8.5] we identified all the ξ_w for which $R_w = B_w$ holds but failed to codify the list in a formal statement. The omission is rectified in Section 5.

The Schubert system lifts to a linear Pfaffian exterior differential system defined on a frame bundle over X [18, Section 4.4]. The lifted system has the advantage that obstructions to flexibility may be identified using Lie algebra cohomology, cf. [18, Sections 4 and 5]. A complete list of the Schubert systems $\mathcal{B}_w$ for which there exist cohomological obstructions to flexibility is given by [18, Theorem 6.1]. It is then shown in [18, Theorem 8.1] that $R_w = B_w$ for each of these rigid systems. In particular, the corresponding Schur system $\mathcal{R}_w$ is rigid.

The first goal of this paper is to prove that these cohomological obstructions are a complete set of obstructions to the Schubert flexibility of ξ_w. That is, in the absence of cohomological obstructions there exist nontrivial, algebraic integrals Y of the Schubert system (Theorem 3.1). As an integral of $\mathcal{B}_w$, the variety Y is necessarily an integral of the Schur system $\mathcal{R}_w$. In particular, $[Y] = r\xi_w$ for some integer $r > 0$.

2.4. **The characterization of Schubert varieties.** This section is a brief review of the characterization of Schubert classes ξ_w by an integer $\mathsf{a}(w) \geq 0$ and a marking $\mathsf{J}(w)$ of the Dynkin diagram. (The marking is equivalent to a choice of simple roots.) For more detail see [18, Section 3.2]. Recall (2.4), and let

$$N_w \;=\; \exp(\mathfrak{n}_w)\,.$$

Then

$$X_w \;:=\; \overline{N_w \cdot o} \;=\; wY_w \tag{2.12}$$

is a Schubert variety of type w . (See [18, Sections 2.3 and 2.4] for more detail.)

Let $\mathbb{1} \in W^{\mathfrak{p}}$ be the identity, and let $w_0 \in W^{\mathfrak{p}}$ be the longest element. The $\mathfrak{g}_0$–module $\mathbf{I}_w$ is trivial if and only if $w \in \{\mathbb{1}, w_0\}$. (The associated Schubert varieties are $X_{\mathbb{1}} = o$ and

$X_{w_0} = X$.) Assume $w \in W^{\mathfrak{p}} \backslash \{\mathbb{1}, w_0\}$. Let $\mathfrak{q}_w \subset \mathfrak{g}_0$ be the stabilizer of the highest weight line $\mathfrak{n}_w \in \mathbb{P}(\bigwedge^{|w|} T_o X)$. Then there is a subset $\mathtt{J}(w) \subset \{1, \ldots, r\} \backslash \{\mathtt{i}\}$ with the property that the Lie algebra $\mathfrak{q}_w$ is given by $\mathfrak{q}_w = \mathfrak{g}_{0,\geq 0}$, where

$$\mathfrak{g}_{k,\ell} \ := \ \{A \in \mathfrak{g}_k \mid [Z_w\,,\, A] = \ell A\} \quad \text{and} \quad Z_w = \sum_{\mathtt{j} \in \mathtt{J}(w)} Z_{\mathtt{j}}\,. \tag{2.13}$$

We call $\mathfrak{g} = \oplus \mathfrak{g}_{k,\ell}$ the $(Z_{\mathtt{i}}, Z_w)$*–bigraded decomposition* of $\mathfrak{g}$. The following is [18, Proposition 3.9].

Proposition 2.14 ([18])**.** *Let* $w \in W^{\mathfrak{p}} \backslash \{\mathbb{1}, w_0\}$*. There exists an integer* $\mathtt{a} = \mathtt{a}(w) \geq 0$ *such that* $\Delta(w) = \{\alpha \in \Delta(\mathfrak{g}_1) \mid \alpha(Z_w) \leq \mathtt{a}\}$*. Equivalently,*

$$\mathfrak{n}_w \ = \ \mathfrak{g}_{-1,0} \ \oplus \ \cdots \ \oplus \ \mathfrak{g}_{-1,-\mathtt{a}}\,. \tag{2.15}$$

Remarks. (1) Since $\xi_w = [X_w]$, and X_w is determined by $\Delta(w)$, the pair $\mathtt{a}(w), \mathtt{J}(w)$ characterizes ξ_w, when $w \in W^{\mathfrak{p}}\{\mathbb{1}, w_0\}$.

(2) By [18, Proposition 3.19], the Schubert variety X_w is smooth if and only if $\mathtt{a}(w) = 0$. For more on the relationship between the integer $\mathtt{a}(w)$ and $\mathrm{Sing}(X_w)$, see [17].

(3) A tableau-esque analog of Proposition 2.14 is given by H. Thomas and A. Yong in [20, Proposition 2.1].

A complete list of the $\mathtt{a}(w)$, $\mathtt{J}(w)$ that occur is given by [18, Corollary 3.17]. For each of the classical, cominuscule varieties, we review the $(\mathtt{a}, \mathtt{J})$–description associated to the Schubert variety X_w in Sections 2.5–2.10. This includes a discussion of the correspondence between the $(\mathtt{a}, \mathtt{J})$–data and the more familiar partition–based descriptions. The values of $(\mathtt{a}, \mathtt{J})$ corresponding to Schubert varieties in the two exceptional cominuscule varieties are given by Figures 1 and 2.

Notation. Given a basis $\{e_1, \ldots, e_n\}$ of a vector space V, let $\{e^1, \ldots, e^n\}$ denote the dual basis of V^*. Set $e^a_b = e_b \otimes e^a \in \mathrm{End}(V)$ for all $1 \leq a, b \leq n$.

Remark 2.16. I follow the notation of [18], with the following exception. In [18], we uniformly write $\mathtt{J} = \{\mathtt{j}_1 < \cdots < \mathtt{j}_\mathtt{p}\}$. In this paper, it is convenient to reorder the $\mathtt{j}_\ell$ in some cases.

2.5. Quadric hypersurfaces $\mathcal{Q}^m \subset \mathbb{P}^{m+1}$.

2.5.1. *Odd dimensional quadrics* $\mathcal{Q}^{2n-1} = B_n/P_1$. Set $m = 2n-1$. There is a bijection between $W^{\mathfrak{p}} \backslash \{\mathbb{1}, w_0\}$ and pairs $(\mathtt{a}, \mathtt{J})$ such that $\mathtt{J} = \{\mathtt{j}\} \subset \{2, \ldots, n\}$ and $\mathtt{a} \in \{0, 1\}$; see [18, Corollary 3.17]. The Schubert variety X_w associated with a fixed $\mathtt{a}$ and $\mathtt{J}$ may be described as follows.

Given a nondegenerate symmetric quadric form $(\cdot, \cdot)$ on $\mathbb{C}^{2n+1}$, fix a basis $\{e_1, \ldots, e_{2n+1}\}$ of $\mathbb{C}^{2n+1}$ so that $(e_k, e_\ell) = (e_{n+k}, e_{n+\ell}) = (e_k, e_{2n+1}) = (e_{n+k}, e_{2n+1}) = 0$, $(e_k, e_{n+\ell}) = \delta_{k\ell}$, for all $1 \leq k, \ell \leq n$, and $(e_{2n+1}, e_{2n+1}) = 1$. The abelian subalgebra $\mathfrak{g}_{-1}$ is spanned by the root vectors

$$\begin{array}{rll} e^1_k - e^{n+k}_{n+1}\,, & \text{with root} & -(\alpha_1 + \cdots + \alpha_{k-1}), \\ e^1_{n+k} - e^k_{n+1}\,, & \text{with root} & -\left(\alpha_1 + \cdots + \alpha_{k-1} + 2(\alpha_k + \cdots + \alpha_n)\right), \\ e^1_{2n+1} - e^{2n+1}_{n+1}\,, & \text{with root} & -(\alpha_1 + \cdots + \alpha_n)\,, \end{array}$$

$2 \le k \le n$.

Set $o = [e_1] \in \mathbb{P}^{2n}$. If $\mathtt{a} = 0$, then

$$\mathfrak{n}_w = \langle e^1_k - e^{n+k}_{n+1} \mid k \le \mathtt{j} \rangle \quad \text{and} \quad X_w = \overline{\exp(\mathfrak{n}_w) \cdot o} = \mathbb{P}\langle e_1, \ldots, e_{\mathtt{j}} \rangle = \mathbb{P}^{\mathtt{j}-1} \,.$$

If $\mathtt{a} = 1$, then

$$\mathfrak{n}_w = \langle e^1_k - e^{n+k}_{n+1} \,,\ e^1_{n+\ell} - e^{\ell}_{n+1} \,,\ e^1_{2n+1} - e^{2n+1}_{n+1} \mid 1 \le k \le n \,,\ \mathtt{j} < \ell \rangle$$

and $X_w = \overline{\exp(\mathfrak{n}_w) \cdot o} = \mathcal{Q}^m \cap \mathbb{P}\langle e_1, \ldots, e_{n+1}, e_{n+\mathtt{j}+1}, \ldots, e_{2n+1} \rangle$.

2.5.2. *Even dimensional quadrics* $\mathcal{Q}^{2n-2} = D_n/P_1$. Set $m = 2n-2$. There is a bijection between $W^{\mathfrak{p}} \backslash \{\mathbb{1}, w_0\}$ and pairs $(\mathtt{a}, \mathtt{J})$ such that either

- ◦ $\mathtt{a} = 0$ and $\mathtt{J} = \{\mathtt{j}\} \subset \{2, \ldots, n\}$ or $\mathtt{J} = \{n-1, n\}$; or
- ◦ $\mathtt{a} = 1$ and $\mathtt{J} = \{\mathtt{j}\} \subset \{2, \ldots, n-2\}$ or $\mathtt{J} = \{n-1, n\}$.

See [18, Corollary 3.17]. The Schubert variety X_w associated with a fixed $\mathtt{a}$ and $\mathtt{J}$ may be described as follows.

Given a nondegenerate symmetric quadric form $(\cdot,\cdot)$ on $\mathbb{C}^{2n}$, fix a basis $\{e_1, \ldots, e_{2n}\}$ of $\mathbb{C}^{2n}$ so that $(e_k, e_\ell) = (e_{n+k}, e_{n+\ell}) = 0$ and $(e_k, e_{n+\ell}) = \delta_{k\ell}$, for all $1 \le k, \ell \le n$. The abelian subalgebra $\mathfrak{g}_{-1}$ is spanned by the root vectors

$$\begin{aligned} e^1_k - e^{n+k}_{n+1} \,, &\quad \text{with root} \quad -(\alpha_1 + \cdots + \alpha_{k-1}), \\ e^1_{n+\ell} - e^{\ell}_{n+1} \,, &\quad \text{with root} \quad -\left(\alpha_1 + \cdots + \alpha_{\ell-1} + 2(\alpha_\ell + \cdots + \alpha_{n-2}) + \alpha_{n-1} + \alpha_n\right), \\ e^1_{2n} - e^{n}_{n+1} \,, &\quad \text{with root} \quad -(\alpha_1 + \cdots + \alpha_{n-2}) - \alpha_n, \end{aligned}$$

with $2 \le k \le n$ and $2 \le \ell \le n-1$.

Set $o = [e_1] \in \mathbb{P}\mathbb{C}^{2n}$. First suppose that $\mathtt{a} = 0$. If $\mathtt{J} = \{\mathtt{j}\}$ with $2 \le \mathtt{j} \le n-2$, then $X_w = \mathbb{P}^{\mathtt{j}-1}$. If $\mathtt{J} = \{n-1\}$ or $\mathtt{J} = \{n\}$, then $X_w = \mathbb{P}^{n-1}$. If $\mathtt{J} = \{n-1, n\}$, then $X_w = \mathbb{P}^{n-2}$.

Next suppose that $\mathtt{a} = 1$. If $\mathtt{J} = \{\mathtt{j}\}$ with $2 \le \mathtt{j} \le n-2$, then $X_w = \mathcal{Q}^m \cap \mathbb{P}\langle e_1, \ldots, e_{n+1}, e_{n+\mathtt{j}+1}, \ldots, e_{2n} \rangle$. If $\mathtt{J} = \{n-1, n\}$, then $X_w = \mathcal{Q}^m \cap \mathbb{P}\langle e_1, \ldots, e_{n+1}, e_{2n} \rangle$.

2.6. **Grassmannians $\mathrm{Gr}(\mathtt{i}, n+1) = A_n/P_{\mathtt{i}}$.** There is a bijection between $W^{\mathfrak{p}} \backslash \{\mathbb{1}, w_0\}$ and pairs $(\mathtt{a}, \mathtt{J})$ such that $\mathtt{J} = \{\mathtt{j}_\mathtt{p}, \ldots, \mathtt{j}_1, \mathtt{k}_1, \ldots, \mathtt{k}_\mathtt{q}\} \subset \{1, \ldots, n\} \backslash \{\mathtt{i}\}$ is ordered so that

$$1 \le \mathtt{j}_\mathtt{p} < \cdots < \mathtt{j}_1 < \mathtt{i} < \mathtt{k}_1 < \cdots < \mathtt{k}_\mathtt{q} \le n \,,$$

and satisfying $\mathtt{p}, \mathtt{q} \in \{\mathtt{a}, \mathtt{a}+1\}$; see [18, Corollary 3.17]. (Beware, these $\mathtt{p}, \mathtt{q}$ do not agree with those of [18], cf. Remark 2.16.) For convenience we set

$$\mathtt{j}_{\mathtt{p}+1} := 0 \,, \quad \mathtt{j}_0 := \mathtt{i} =: \mathtt{k}_0 \,, \quad \mathtt{k}_{\mathtt{q}+1} := n+1 \,.$$

Fix a basis $\{e_1, \ldots, e_{n+1}\}$ of $\mathbb{C}^{n+1}$. The abelian subalgebra $\mathfrak{g}_{-1}$ is spanned by the root vectors $\{e^k_\ell \mid 1 \le k \le \mathtt{i} < \ell \le n+1\}$; the corresponding roots are $-(\alpha_k + \cdots + \alpha_{\ell-1})$. Define a filtration $\mathtt{F}_{\mathtt{a}+1} \subset \mathtt{F}_\mathtt{a} \subset \cdots \subset \mathtt{F}_1 \subset \mathtt{F}_0$ of $\mathbb{C}^{n+1}$ by

$$\mathtt{F}_\ell = \langle e_1, e_2, \ldots, e_{\mathtt{j}_\ell} \,,\ e_{\mathtt{i}+1}, e_{\mathtt{i}+2}, \ldots, e_{\mathtt{k}_m} \rangle \quad \text{with} \quad \ell + m = \mathtt{a} + 1 \,. \tag{2.17a}$$

Set $o = [e_1 \wedge \cdots \wedge e_{\mathtt{i}}] \in \mathbb{P}(\bigwedge^{\mathtt{i}} \mathbb{C}^{n+1})$. Then

$$X_w = \{E \in \mathrm{Gr}(\mathtt{i}, n+1) \mid \dim(E \cap \mathtt{F}_\ell) \ge \mathtt{j}_\ell \,,\ 0 \le \ell \le \mathtt{a}+1\} \,. \tag{2.17b}$$

Example 2.18. Consider $X = \mathrm{Gr}(5,13) \simeq A_{12}/P_5$. The marking $\mathtt{J} = \{2,3,7,9,12\}$ and integer $\mathtt{a} = 2$ define the filtration $\mathtt{F}_3 \subset \mathtt{F}_2 \subset \mathtt{F}_1 \subset \mathtt{F}_0$ as

$$\begin{aligned} \mathtt{F}_3 &= \langle 0 \rangle\,, \quad \mathtt{F}_2 \;=\; \langle e_1, e_2\,,\, e_6, e_7 \rangle\,,\\ \mathtt{F}_1 &= \langle e_1, e_2, e_3\,,\, e_6, e_7, e_8, e_9 \rangle\,, \quad \mathtt{F}_0 \;=\; \langle e_1, \ldots, e_5\,,\, e_6, \ldots e_{12} \rangle\,. \end{aligned}$$

The associated Schubert variety is the set of all $E \in \mathrm{Gr}(5,13)$ such that

$$\dim(E \cap \mathtt{F}_3) \geq 0\,, \quad \dim(E \cap \mathtt{F}_2) \geq 2\,, \quad \dim(E \cap \mathtt{F}_1) \geq 3\,, \quad \dim(E \cap \mathtt{F}_0) \geq 5\,.$$

Example 2.19. Consider $X = \mathrm{Gr}(6,11) \simeq A_{10}/P_6$. The marking $\mathtt{J} = \{1,3,4,7,9\}$ and integer $\mathtt{a} = 2$ define the filtration $\mathtt{F}_3 \subset \mathtt{F}_2 \subset \mathtt{F}_1 \subset \mathtt{F}_0$ as

$$\begin{aligned} \mathtt{F}_3 &= \langle e_1 \rangle\,, \quad \mathtt{F}_2 \;=\; \langle e_1, e_2, e_3\,,\, e_7 \rangle\,,\\ \mathtt{F}_1 &= \langle e_1, e_2, e_3, e_4\,,\, e_7, e_8, e_9 \rangle\,, \quad \mathtt{F}_0 \;=\; \mathbb{C}^{11}\,. \end{aligned}$$

The associated Schubert variety is the set of all $E \in \mathrm{Gr}(6,11)$ such that

$$\dim(E \cap \mathtt{F}_3) \geq 1\,, \quad \dim(E \cap \mathtt{F}_2) \geq 3\,, \quad \dim(E \cap \mathtt{F}_1) \geq 4\,, \quad \dim(E \cap \mathtt{F}_0) \geq 6\,.$$

2.7. **Partitions versus $(\mathtt{a}, \mathtt{J})$ in Grassmannians.** Complete the filtration $\mathtt{F}_\bullet$ of (2.17a) to a full flag

$$0 \subset F^1 \subset F^2 \subset \cdots \subset F^{n+1}\,, \quad \dim F^d = d\,. \tag{2.20}$$

Note that

$$\mathtt{F}_\ell \;=\; F^{\mathtt{j}_\ell + \mathtt{k}_m - \mathtt{i}}\,, \quad \text{where} \quad \ell + m = \mathtt{a} + 1\,.$$

It is well-known that Schubert varieties in $\mathrm{Gr}(\mathtt{i}, n+1)$ are indexed by partitions

$$\lambda \;=\; (\lambda_1, \ldots, \lambda_{\mathtt{i}}) \in \mathbb{Z}^{\mathtt{i}} \quad \text{such that} \quad 1 \leq \lambda_1 < \lambda_2 < \cdots < \lambda_{\mathtt{i}} \leq n+1\,. \tag{2.21}$$

The corresponding Schubert variety is

$$X_\lambda \;:=\; \{E \in \mathrm{Gr}(\mathtt{i}, n+1) \mid \dim(E \cap F^{\lambda_k}) \geq k\,,\ 1 \leq k \leq \mathtt{i}\}\,. \tag{2.22}$$

We may determine the $\mathtt{a}(\lambda)$ and $\mathtt{J}(\lambda)$ values associated to X_λ (cf. Proposition 2.14) as follows. Note that, if $\lambda_{k+1} = \lambda_k + 1$, then the condition $\dim(E \cap F^{\lambda_k}) \geq k$ is redundant; it is implied by $\dim(E \cap F^{\lambda_{k+1}}) \geq k+1$. To remove the redundancies, decompose $\lambda = \mu^{\mathtt{p}} \cdots \mu^1 \mu^0$ into maximal blocks of consecutive integers. For example, if $\lambda = (2,3,4,7,8,12)$, then $\mu^2 = (2,3,4)$, $\mu^1 = (7,8)$ and $\mu^0 = (12)$. Define

$$\mathtt{j}_\ell(\lambda) \;=\; |\mu^{\mathtt{p}} \cdots \mu^\ell| \tag{2.23}$$

to be the length of the sub-partition $\mu^{\mathtt{p}} \cdots \mu^\ell$. (In all cases, $\mathtt{j}_0 = |\lambda| = \mathtt{i}$.) In the preceeding example, $\mathtt{j}_2 = 3$, $\mathtt{j}_1 = 5$ and $\mathtt{j}_0 = 6$. Note that $\lambda_{\mathtt{j}_\ell}$ is the last entry in the block μ^ℓ. That is,

$$\{\lambda_{\mathtt{j}_{\mathtt{p}}}, \ldots, \lambda_{\mathtt{j}_1}\} \;=\; \{\lambda_k \in \lambda \mid \lambda_k - \lambda_{k+1} > 1\,,\ 1 \leq k < \mathtt{i}\}\,. \tag{2.24}$$

The redundancy-free formulation of (2.22) is

$$X_\lambda \;=\; \{E \in \mathrm{Gr}(\mathtt{i}, n+1) \mid \dim(E \cap F^{\lambda_{\mathtt{j}_\ell}}) \geq \mathtt{j}_\ell\,,\ 1 \leq \ell \leq \mathtt{p}\}\,.$$

In particular,

$$F^{\lambda_{\mathtt{j}_\ell}} \;=\; \mathtt{F}_\ell\,.$$

This yields the following reformulation of [18, Proposition 3.30].

Lemma 2.25 ([18])**.** *Let* $\lambda = (\lambda_1,\dots,\lambda_{\mathtt{i}})$ *be a partition satisfying* (2.21)*, and let* $\lambda = \mu^{\mathtt{p}}\cdots\mu^1\mu^0$ *be the decomposition of* λ *into maximal blocks of consecutive integers. The values* $\mathtt{a}(\lambda)$ *and* $\mathtt{J}(\lambda) = \{\mathtt{j}_{\mathtt{p}},\dots,\mathtt{j}_1,\mathtt{k}_1,\dots,\mathtt{k}_{\mathtt{q}}\}$ *are given by* (2.23)*,*

$$\{\mathtt{k}_1,\dots,\mathtt{k}_{\mathtt{q}}\} \;=\; \{\mathtt{i}-\mathtt{j}_{\mathtt{p}}+\lambda_{\mathtt{j}_{\mathtt{p}}},\dots,\ \mathtt{i}-\mathtt{j}_1+\lambda_{\mathtt{j}_1}\,,\ \lambda_{\mathtt{i}}\}\backslash\{\mathtt{i},n+1\}$$

and

$$\mathtt{a} \;=\; \left\{\begin{array}{ll} \mathtt{p} & \text{if } \lambda_1>1 \\ \mathtt{p}-1 & \text{if } \lambda_1 = 1 \end{array}\right\} \;=\; \left\{\begin{array}{ll} \mathtt{q}\,, & \text{if } \lambda_{\mathtt{i}} = n+1 \\ \mathtt{q}-1\,, & \text{if } \lambda_{\mathtt{i}} < n+1\,. \end{array}\right.$$

Conversely, given $(\mathtt{a},\mathtt{J})$*, the associated partition* $\lambda = \mu^{\mathtt{p}}\cdots\mu^1\mu^0$ *is given by*

$$\mu^\ell \;=\; (\mathtt{j}_{\ell+1}+\mathtt{k}_m-\mathtt{i}+1\,,\dots,\ \mathtt{j}_\ell+\mathtt{k}_m-\mathtt{i})\,,$$

with $\ell+m = \mathtt{a}+1$*.*

Example. In Example 2.18, we have $\lambda = (3,4,7,11,12)$; and in Example 2.19 we have $\lambda = (1,3,4,7,10,11)$.

Remark. Together [18, Proposition 3.30] and [1, Theorem 9.3.1] imply that the integer $\mathtt{a}(w)$ is the number of irreducible components of $\mathrm{Sing}(X_w)$. See [17] for a uniform (type independent) description of the singular loci of cominuscule Schubert varieties.

2.8. **Lagrangian Grassmannians $\mathbf{LG}(n,2n) = C_n/P_n$.** There exists a bijection between $W^{\mathtt{p}}\backslash\{\mathbb{1},w_0\}$ and pairs $\mathtt{a}\geq 0$ and $\mathtt{J} = \{\mathtt{j}_{\mathtt{p}},\dots,\mathtt{j}_1\}\subset\{1,\dots,n-1\}$ satisfying

$$1 \;\leq\; \mathtt{j}_{\mathtt{p}} < \cdots < \mathtt{j}_1 \;\leq\; n-1$$

and $|\mathtt{J}| = \mathtt{p}\in\{\mathtt{a},\mathtt{a}+1\}$; see [18, Corollary 3.17]. (Note that these $\mathtt{j}_\ell$ have the opposite order of those in [18].) For convenience we set

$$\mathtt{j}_{\mathtt{p}+1} := 0\,,\quad \mathtt{j}_0 := n\,.$$

To describe the corresponding Schubert variety X_w, fix a nondegenerate skew-symmetric bilinear form $(\cdot,\cdot)$ on $\mathbb{C}^{2n}$, and basis $\{e_1,\dots,e_{2n}\}$ of $\mathbb{C}^{2n}$ satisfying $(e_a,e_b) = 0 = (e_{n+a},e_{n+b})$ and $1 = (e_a,e_{n+b}) = \delta_{ab}$ for all $1\leq a,b\leq n$. Then $\mathrm{LG}(n,2n)$ is the C_n–orbit of $\langle e_1,\dots,e_n\rangle$. The abelian subalgebra $\mathfrak{g}_{-1}$ (which may be identified with n-by-n symmetric matrices) is spanned by the root vectors $\{e^j_{n+k}+e^k_{n+j}\mid 1\leq j\leq k\leq n\}$, with roots $-(\alpha_j+\cdots+\alpha_{k-1})-2(\alpha_k+\cdots+\alpha_n)$. Define a filtration $\mathtt{F}_{\mathtt{p}}\subset\cdots\subset\mathtt{F}_1\subset\mathtt{F}_0$ of $\mathbb{C}^{2n}$ by

$$\mathtt{F}_\ell \;=\; \langle e_1,\dots,e_{\mathtt{j}_\ell}\,,\ e_{n+\mathtt{j}_m+1},\dots,e_{2n}\rangle \quad\text{with}\quad \ell+m = \mathtt{a}+1\,. \tag{2.26a}$$

Set $o = [e_1\wedge\cdots\wedge e_n]\in\mathbb{P}(\bigwedge^n\mathbb{C}^{2n})$. Then

$$X_w \;=\; \{E\in\mathrm{LG}(n,2n)\mid \dim(E\cap\mathtt{F}_\ell)\geq\mathtt{j}_\ell\,,\ \forall\ 0\leq\ell\leq\mathtt{p}\}\,. \tag{2.26b}$$

Example 2.27. Consider $X = \mathrm{LG}(5,10)\simeq C_5/P_5$. The marking $\mathtt{J} = \{1,2,4\}$ and integer $\mathtt{a} = 3$ define the filtration $\mathtt{F}_3\subset\mathtt{F}_2\subset\mathtt{F}_1\subset\mathtt{F}_0$ of $\mathbb{C}^{10}$ as

$$\begin{aligned} \mathtt{F}_3 \;&=\; \langle e_1\,,\,e_{10}\rangle\,,\quad \mathtt{F}_2 \;=\; \langle e_1,e_2\,,\,e_8,e_9,e_{10}\rangle\,,\\ \mathtt{F}_1 \;&=\; \langle e_1,\dots,e_4\,,\,e_7,\dots,e_{10}\rangle\,,\quad \mathtt{F}_0 \;=\; \mathbb{C}^{10}\,. \end{aligned}$$

The associated Schubert variety is the set of all $E\in\mathrm{LG}(5,10)$ such that

$$\dim(E\cap\mathtt{F}_3)\geq 1\,,\quad \dim(E\cap\mathtt{F}_2)\geq 2\,,\quad \dim(E\cap\mathtt{F}_1)\geq 4\,.$$

2.9. **Partitions versus $(\mathsf{a},\mathtt{J})$ in Lagrangian grassmannians.** Complete the filtration $\mathtt{F}_\bullet$ of (2.26a) to a full isotropic flag $F^\bullet$; that is,

$$(F^k, F^{2n-k}) \;=\; 0 \ \text{ for all } 1 \le k \le n\,, \quad \text{and} \quad \mathtt{F}_\ell \;=\; F^{\mathtt{j}_\ell + n - \mathtt{j}_m} \ \text{ where } \ \ell + m = \mathsf{a}+1\,. \tag{2.28}$$

It is well-known that Schubert varieties in $\mathrm{LG}(n,2n)$ are indexed by partitions $\lambda = (\lambda_1, \lambda_2, \ldots, \lambda_n)$ such that

$$1 \le \lambda_1 < \lambda_2 < \cdots < \lambda_n \le 2n\,, \quad \text{and} \tag{2.29a}$$

$$\lambda_i \in \lambda \quad \text{if and only if} \quad 2n+1-\lambda_i \not\in \lambda\,. \tag{2.29b}$$

The corresponding Schubert variety is

$$X_\lambda \;=\; \{E \in \mathrm{LG}(n,2n) \mid \dim(E \cap F^{\lambda_k}) \ge k\,,\ 1 \le k \le n\}\,. \tag{2.30}$$

We may determine the values $\mathsf{a}(\lambda)$ and $\mathtt{J}(\lambda)$ associated to X_λ (cf. Proposition 2.14) as follows. As was the case in Section 2.6, the definition of X_λ is redundant. Again, to remove the redundancies, decompose $\lambda = \mu^{\mathsf{p}} \cdots \mu^1 \mu^0$ into maximal blocks of consecutive integers. Let $\mathtt{j}_\ell(\lambda) = |\mu^{\mathsf{p}} \cdots \mu^\ell|$ be the length of the sub-partition $\mu^{\mathsf{p}} \cdots \mu^s$. (In all cases, $\mathtt{j}_0 = n$.) Then the redundancy-free formulation is

$$X_\lambda \;=\; \{E \in \mathrm{LG}(n,2n) \mid \dim(E \cap F^{\lambda_{\mathtt{j}_\ell}}) \ge \mathtt{j}_\ell\,,\ 1 \le \ell \le \mathsf{p}\}\,.$$

A comparison of this with (2.26b) yields $\mathtt{J} = \{\mathtt{j}_{\mathsf{p}}(\lambda), \ldots, \mathtt{j}_1(\lambda)\}$ and

$$\mathtt{F}_\ell \;=\; F^{\lambda_{\mathtt{j}_\ell}}\,,$$

so that $\lambda_{\mathtt{j}_\ell} = \dim \mathtt{F}_\ell = \mathtt{j}_\ell + n - \mathtt{j}_m$, for $\ell + m = \mathsf{a}+1$; note that $\lambda_{\mathtt{j}_\ell} + \lambda_{\mathtt{j}_m} = 2n$. In order to determine the value of a, we consider two cases:

- First, suppose that $\lambda_1 > 1$. By (2.29b), this is equivalent to $\lambda_n = \lambda_{\mathtt{j}_0} = 2n$. Additionally, (2.29b) yields $\lambda_{\mathtt{j}_\ell} + \lambda_{\mathtt{j}_m} = 2n$ when $\ell + m = \mathsf{p}+1$. Therefore, $\mathsf{a} = \mathsf{p}$.
- Second, suppose that $\lambda_1 = 1$, equivalently, $\lambda_n = \lambda_{\mathtt{j}_0} < 2n$. Equation (2.29b) implies $\lambda_{\mathtt{j}_\ell} + \lambda_{\mathtt{j}_m} = 2n$ when $\ell + m = \mathsf{p}$. Thus, $\mathsf{p} = \mathsf{a}+1$.

The preceding discussion may be summarized as follows.

Lemma 2.31. *Let $\lambda = (\lambda_1, \ldots, \lambda_n)$ be a partition satisfying* (2.29). *Let $\lambda = \mu^{\mathsf{p}} \cdots \mu^1 \mu^0$ be a decomposition of λ into $\mathsf{p}+1$ maximal blocks of consecutive integers. Then $\mathtt{J}(\lambda) = \{\mathtt{j}_{\mathsf{p}}(\lambda), \ldots, \mathtt{j}_1(\lambda)\}$ is given by* (2.23), *and*

$$\mathsf{a}(\lambda) \;=\; \begin{cases} \mathsf{p}-1 & \text{if } \lambda_1 = 1 \\ \mathsf{p} & \text{if } \lambda_1 > 1\,. \end{cases}$$

Conversely, given a and $\mathtt{J} = \{\mathtt{j}_{\mathsf{p}}, \cdots, \mathtt{j}_1\}$ we construct $\lambda(\mathsf{a},\mathtt{J}) = \mu^{\mathsf{p}}(\mathsf{a},\mathtt{J}) \cdots \mu^0(\mathsf{a},\mathtt{J})$ by

$$\mu^\ell(\mathsf{a},\mathtt{J}) \;=\; (n+1+\mathtt{j}_{\ell+1} - \mathtt{j}_m\,,\ \ldots\,,\ n + \mathtt{j}_\ell - \mathtt{j}_m)\,, \quad \textit{with} \quad \ell + m = \mathsf{a}+1\,. \tag{2.32}$$

Example. Consider $n = 15$ and $\mathtt{J} = \{3,7,9,13\}$. Since $|\mathtt{J}| = 4$, we have either $\mathsf{a} = 3$ or $\mathsf{a} = 4$. If $\mathsf{a} = 3$, then the partition is $\lambda = (1,2,3)(6,7,8,9)(14,15)(18,19,20,21,22)(26,27)$. If $\mathsf{a} = 4$, then $\lambda = (3,4,5)(10,11,12,13)(16,17)(22,23,24,25)(29,30)$.

2.10. **Spinor varieties $\mathcal{S}_n = D_n/P_n$.** Given $\mathsf{a} = \mathsf{a}(w)$ and $\mathtt{J} = \mathtt{J}(w)$, note that

$$\alpha_{n-1}(Z_w) = 0 \ \text{ if } n-1 \notin \mathtt{J}\,, \qquad \text{and} \qquad \alpha_{n-1}(Z_w) = 1 \ \text{ if } n-1 \in \mathtt{J}\,.$$

Define

$$\mathsf{r} \ = \ \left\lceil \tfrac{1}{2}\left(\mathsf{a} + \alpha_{n-1}(Z_w)\right)\right\rceil \ = \ \begin{cases} \lceil \mathsf{a}/2 \rceil & \text{if } n-1 \notin \mathtt{J}\,, \\ \lfloor \mathsf{a}/2 \rfloor + 1 & \text{if } n-1 \in \mathtt{J}\,. \end{cases} \tag{2.33}$$

There exists a bijection between $W^{\mathfrak{p}}\backslash\{\mathbb{1}, w_0\}$, and pairs $\mathsf{a} \geq 0$ and $\mathtt{J} = \{\mathtt{j}_{\mathsf{p}}, \ldots, \mathtt{j}_1\} \subset \{1, \ldots, n-1\}$, ordered so that

$$1 \ \leq \ \mathtt{j}_{\mathsf{p}} < \cdots < \mathtt{j}_1 \ \leq \ n-1\,, \tag{2.34a}$$

and satisfying

$$\mathsf{p} - \alpha_{n-1}(Z_w) \in \{\mathsf{a}, \mathsf{a}+1\}\,, \quad \text{and} \quad 2 \leq \mathtt{j}_{\mathsf{r}} - \mathtt{j}_{\mathsf{r}+1} \ \text{ when } \ \mathsf{r} > \alpha_{n-1}(Z_w)\,; \tag{2.34b}$$

see [18, Corollary 3.17]. (Note that these $\mathtt{j}_\ell$ have the opposite order of those in [18].) For convenience we set

$$\mathtt{j}_{\mathsf{p}+1} := 0\,, \quad \mathtt{j}_0 := n\,.$$

To describe the corresponding Schubert variety X_w, fix a nondegenerate symmetric bilinear form $(\cdot,\cdot)$ on $\mathbb{C}^{2n}$ and basis $\{e_1, \ldots, e_{2n}\}$ of $\mathbb{C}^{2n}$ satisfying $(e_a, e_b) = 0 = (e_{n+a}, e_{n+b})$ and $1 = (e_a, e_{n+b}) = \delta_{ab}$ for all $1 \leq a, b \leq n$. Our convention is that $\mathcal{S}_n$ is the D_n–orbit of $\langle e_1, \ldots, e_n\rangle$. The abelian subalgebra $\mathfrak{g}_{-1}$ (which may be identified with n-by-n skew-symmetric matrices) is spanned by root vectors $\{e^j_{n+k} - e^k_{n+j} \mid 1 \leq j < k \leq n\}$. The corresponding roots are $-(\alpha_j + \cdots + \alpha_{n-2}) - \alpha_n$, if $k = n$; $-(\alpha_j + \cdots + \alpha_n)$, if $k = n-1$; and $-(\alpha_j + \cdots + \alpha_{k-1}) - 2(\alpha_k + \cdots + \alpha_{n-2}) - \alpha_{n-1} - \alpha_n$, if $k < n-1$. Define a filtration $\mathtt{F}_{\mathsf{p}} \subset \cdots \subset \mathtt{F}_1 \subset \mathtt{F}_0$ of $\mathbb{C}^{2n}$ by

$$\mathtt{F}_\ell \ = \ \langle e_1, \ldots, e_{\mathtt{j}_\ell}\,, e_{n+\mathtt{j}_m+1}, \ldots, e_{2n}\rangle \tag{2.35a}$$

with

$$\ell + m \ = \ \left\{ \begin{array}{ll} \mathsf{a}+1 & \text{if } n-1 \notin \mathtt{J}\,, \\ \mathsf{a}+2 & \text{if } n-1 \in \mathtt{J} \end{array} \right\} \ = \ \mathsf{a} + 1 + \alpha_{n-1}(Z_w)\,. \tag{2.35b}$$

Set $o = [e_1 \wedge \cdots \wedge e_n] \in \mathbb{P}(\bigwedge^n \mathbb{C}^{2n})$. Then

$$X_w \ = \ \{E \in \mathcal{S}_n \mid \dim(E \cap \mathtt{F}_\ell) \geq \mathtt{j}_\ell\,, \ \forall\ 0 \leq \ell \leq \mathsf{p}\}\ . \tag{2.35c}$$

2.11. **Partitions versus $(\mathsf{a}, \mathtt{J})$ in $\mathcal{S}_n$.** Complete the filtration $\mathtt{F}_\bullet$ of (2.35a) to a full isotropic flag $F^\bullet$; that is, we have (2.28), where the nondegenerate $(\cdot,\cdot)$ is now taken to be symmetric. The partition description for the Schubert varieties in the spinor variety is very similar to those in the Lagrangian grassmannian. Begin with a partition λ satisfying (2.29). Let $\mathtt{J}(\lambda)$ be given by (2.23), define

$$\mathsf{a}(\lambda) \ = \ \begin{cases} \mathsf{p}-1 & \text{if } \lambda_1 = 1 \text{ and } \mathtt{j}_1(\lambda) < n-1\,, \\ \mathsf{p} & \text{if } \lambda_1 > 1 \text{ and } \mathtt{j}_1(\lambda) < n-1\,, \\ \mathsf{p}-2 & \text{if } \lambda_1 = 1 \text{ and } \mathtt{j}_1(\lambda) = n-1\,, \\ \mathsf{p}-1 & \text{if } \lambda_1 > 1 \text{ and } \mathtt{j}_1(\lambda) = n-1\,, \end{cases} \tag{2.36}$$

and let $\mathsf{r}(\lambda)$ be given by (2.33). Then the Schubert varieties of $\mathcal{S}_n$ are indexed by partitions $\lambda = (\lambda_1, \lambda_2, \ldots, \lambda_n)$ satisfying (2.29) and

$$2 \ \leq \ \mathsf{j}_\mathsf{r} - \mathsf{j}_{\mathsf{r}+1} \ = \ |\mu^\mathsf{r}| \,. \tag{2.37}$$

The corresponding Schubert variety is given by

$$X_\lambda \ = \ \{E \in \mathcal{S}_n \mid \dim(E \cap F^{\lambda_k}) \geq k \,, \ 1 \leq k \leq n\} \,. \tag{2.38}$$

Example. Consider $n = 15$ and $\mathsf{J} = \{3, 6, 7, 11, 12\}$. The constraint $2 \geq \mathsf{j}_\mathsf{r} - \mathsf{j}_{\mathsf{r}+1}$ implies $\mathsf{r} \neq 1, 3$. So $\mathsf{a} \neq 1, 2, 5, 6$. Since $\mathsf{p} = 5$ and $\mathsf{j}_1 = 12 < n - 1 = 14$, this forces $\mathsf{a} = 4$. The corresponding partition is $\lambda = (1, 2, 3)(7, 8, 9)(11)(16, 17, 18, 19)(21)(25, 26, 27)$.

Example. Consider $n = 15$ and $\mathsf{J} = \{1, 5, 8, 10, 14\}$. Then $\mathsf{a} = 3$ or $\mathsf{a} = 4$. If $\mathsf{a} = 3$, then $\lambda = (1)(3, 4, 5, 6)(11, 12, 13)(16, 17)(21, 22, 23, 24)(29)$. If $\mathsf{a} = 4$, then

$$\lambda = (2)(7, 8, 9, 10)(13, 14, 15)(19, 20)(25, 26, 27, 28)(30) \,.$$

In analogy with Lemma 2.31, we have the following relationship between the partition and (a, J) descriptions (cf. Proposition 2.14) of Schubert varieties in $\mathcal{S}_n$.

Lemma 2.39. *Schubert varieties in* $\mathcal{S}_n = D_n/P_n$ *are indexed by partitions*

$$\lambda = (\lambda_1, \ldots, \lambda_n) = \mu^\mathsf{p} \cdots \mu^1 \mu^0$$

satisfying (2.29) *and* (2.37)*; the corresponding Schubert variety is given by* (2.38)*. The set* $\mathsf{J}(\lambda) = \{\mathsf{j}_\mathsf{p}(\lambda), \ldots, \mathsf{j}_1(\lambda)\}$ *is given by* (2.23)*, and* $\mathsf{a}(\lambda)$ *is given by* (2.36)*. Conversely, given* a *and* J*, the associated partition is given by* (2.32) *with* $\ell + m = \mathsf{a} + 1 + \alpha_{n-1}(Z_w)$*.*

Remark. Variants of Lemmas 2.31 and 2.39 were also obtained by D. The [19].

2.12. **The exceptional cases.** Figures 1 and 2 (pages 13 and 14) are respectively the Hasse diagrams $W^\mathfrak{p}$ of the Cayley plane E_6/P_6 and Freudenthal variety E_7/P_7. Each node represents a Schubert class ξ_w and is labeled with the corresponding $\mathsf{a}(w) : \mathsf{J}(w)$ values, which were obtained with the assistance of [16]. The height of the node indicates the dimension of X_w; in particular, the lowest node $o \in X$ is at height zero. Two nodes are connected if the Schubert variety associated with the lower node is a divisor of the Schubert variety associated with the higher node.

2.13. **Divisors.** In this section we establish some lemmas that will be used in Section 3. Recall the $(Z_\mathtt{i}, Z_w)$–bigraded decomposition (2.13) of $\mathfrak{g}$, and note that $\mathfrak{g}_{0,0}$ is a reductive subalgebra of $\mathfrak{g}$.

Definition 2.40. Given a representation U of $\mathfrak{g}_{0,0}$, let $\Pi(U)$ denote the set of highest weights.

Given $\alpha \in \Delta^+$, let $r_\alpha \in W$ denote the corresponding reflection.

Lemma 2.41. *Let* $\alpha \in \Delta(w)$*. Then* $\Delta(w)\backslash\{\alpha\} = \Delta(w')$ *for some* $w' \in W^\mathfrak{p}$ *if and only if* $\alpha \in \Pi(\mathfrak{g}_{1,\mathsf{a}})$*. In this case* $w' = r_\alpha w$*.*

Proof. By Remark 2.5, there exists $w' \in W^\mathfrak{p}$ such that $\Delta(w)\backslash\{\alpha\} = \Delta(w')$ if and only if $\Phi = \{\alpha\} \cup (\Delta^+\backslash\Delta(w))$ is closed. So, it suffices to show that $\alpha \in \Pi(\mathfrak{g}_{1,\mathsf{a}})$ if and only if Φ is closed.

Assume $\alpha \in \Pi(\mathfrak{g}_{1,\mathsf{a}})$. Let $\mu, \nu \in \Phi$, and suppose that $\mu + \nu \in \Delta$. To see that Φ is closed, we must show that $\mu + \nu \in \Phi$. First, note that either $\mu \neq \alpha$ or $\nu \neq \alpha$, as $2\alpha \notin \Delta$. Second, if



Figure 1. Hasse diagram of E_6/P_6, each node labeled with the $\mathtt{a} : \mathtt{J}$ values. A node is circled (at least once) if $B_w = R_w$ (Section 5.5), and twice if Schubert the system is rigid (Theorem 4.24).



FIGURE 2. Hasse diagram of E_7/P_7, each node labeled with the $\mathtt{a} : \mathtt{J}$ values. A node is circled (at least once) if $B_w = R_w$ (Section 5.5), and twice if Schubert the system is rigid (Theorem 4.24).

both $\mu, \nu \in \Delta^+ \backslash \Delta(w)$, then $\mu + \nu \in \Delta^+ \backslash \Delta(w) \subset \Phi$, by Remark 2.5. It remains to consider the case that $\mu = \alpha$ and $\nu \in \Delta^+ \backslash \Delta(w)$. Remark 2.3, and the assumption that $\mu + \nu$ is a root, force $\nu \in \Delta^+(\mathfrak{g}_0)$. Note that $\nu(Z_w) \geq 0$. If $\nu(Z_w) = 0$, then ν is a positive root of $\mathfrak{g}_{0,0}$, and the hypothesis that α is a highest $\mathfrak{g}_{0,0}$–weight implies $\mu + \nu \notin \Delta$. If $\nu(Z_w) > 0$, then $(\mu + \nu)(Z_w) > \mathsf{a}$. By Proposition 2.14, we have $\Delta^+ \backslash \Delta(w) = \Delta^+(\mathfrak{g}_0) \cup \Delta(\mathfrak{g}_{1,>\mathsf{a}})$; in particular, $\mu + \nu \in \Delta^+ \backslash \Delta(w) \subset \Phi$.

Conversely, suppose that Φ is closed. To see that α is a highest $\mathfrak{g}_{0,0}$–weight, it suffices to show that $\alpha + \beta \notin \Delta$ for all $\beta \in \Sigma(\mathfrak{g}_{0,0})$. Note that $(\alpha + \beta)(Z_w) = \mathsf{a} + 0$. So, if $\alpha + \beta \in \Delta$, then $\alpha + \beta \in \Delta(\mathfrak{g}_{1,\mathsf{a}})$. By Proposition 2.14, $\Delta(\mathfrak{g}_{1,\mathsf{a}}) \subset \Delta(w)$. Therefore, $\alpha + \beta \in \Delta^+ \backslash \Phi$; since $\alpha, \beta \in \Phi$, this contradicts the assumption that Φ is closed.

It remains to show that $w' = r_\alpha w$. This is [5, Proposition 3.2.15(3)]. □

A similar argument yields the complimentary

Lemma 2.42. *Let $\alpha \in \Delta(\mathfrak{g}_1) \backslash \Delta(w)$. Then $\Delta(w) \sqcup \{\alpha\} = \Delta(w')$ for some $w' \in W^{\mathfrak{p}}$ if and only if $-\alpha \in \Pi(\mathfrak{g}_{-1,-\mathsf{a}-1})$. (Equivalently, α is a lowest $\mathfrak{g}_{0,0}$–weight for $\mathfrak{g}_{1,\mathsf{a}+1}$.) In this case $w' = r_\alpha w$.*

By (2.4) and (2.12), $X_{w'} \subset X_w$ if and only if $\Delta(w') \subset \Delta(w)$. Whence Lemmas 2.41 and 2.42 yield

Corollary 2.43. *Fix a Schubert variety X_w.*

(a) *The number of Schubert divisors in X_w is $|\Pi(\mathfrak{g}_{1,\mathsf{a}})|$.*
(b) *The number of Schubert varieties containing X_w as a divisor is $|\Pi(\mathfrak{g}_{1,\mathsf{a}+1})|$.*

3. Flexibility

In [18, Section 4] we saw that a Lie algebra cohomology $H^1(\mathfrak{n}_w, \mathfrak{g}_w^\perp)$ group is naturally associated with the Schubert system $\mathcal{B}_w$. The cohomology group contains two distinguished subspaces $H^1_{1,\mathsf{a}-1}$ and $H^1_{2,2\mathsf{a}-1}$. When the subspaces are trivial, the Schubert system is necessarily rigid, [18, Theorem 5.38]. The two subspaces are trivial precisely when two representation theoretic conditions, $\mathsf{H}_1 = \mathsf{H}_1(w)$ and $\mathsf{H}_2 = \mathsf{H}_2(w)$, are satisfied (Definition 3.4). When both conditions hold, we say that H_+ is satisfied, and *there exist cohomological obstructions to flexibility.* Theorem 3.1 asserts that in the absence of these cohomological obstructions, the Schubert system is flexible. (In fact, it asserts the stronger statement that ξ_w is Schubert flexible; see Remark 2.7.)

Theorem 3.1. *Let $X = G/P$ be a cominuscule variety, and fix $w \in W^{\mathfrak{p}} \backslash \{\mathbb{1}, w_0\}$. The Schubert system $\mathcal{B}_w$ is rigid if and only if H_+ is satisfied. Moreover, when H_+ fails, there exist nontrivial algebraic integrals; in particular, the Schubert class ξ_w is Schubert rigid if and only if H_+ holds.*

Remark (Geometric interpretation). Recall (Section 2.3) that an irreducible variety $Y \subset X$ is an integral of the Schubert system $\mathcal{B}_w$ if and only if at each smooth point $y \in Y$ there exists a Schubert variety $Y' = Y'(y)$ of type w, such that y is also a smooth point of Y', and $T_y Y = T_y Y'$. Equivalently, if $\mathcal{Y}$ denotes the canonical lift of the smooth locus $Y_0 \subset Y$ to $\mathrm{Gr}(|w|, TX)$, then $\mathcal{Y}$ and $\mathcal{Y}'$ intersect at $\tilde{y} = T_y Y \in \mathrm{Gr}(|w|, TX)$. If the condition H_1 holds, then $\mathcal{Y}$ and $\mathcal{Y}'$ are necessarily tangent at $\tilde{y}$; and when both conditions H_1 and H_2 hold, $\mathcal{Y}$ and $\mathcal{Y}'$ agree to second-order at $\tilde{y}$.

Corollary 3.2. *Let $w \in W^{\mathfrak{p}} \backslash \{\mathbb{1}, w_0\}$. The Schur $\mathcal{R}_w$ system is rigid if and only if H_+ is satisfied. Moreover, when H_+ fails, there exist nontrivial algebraic integrals; in particular, the Schubert class ξ_w is Schur rigid if and only if H_+ holds.*

Proof. The corollary is a direct consequence of Proposition 2.10, Theorem 3.1 and [18, Theorem 8.1]. □

Remark. A complete list of the Schubert varieties (in an irreducible cominuscule X) satisfying H_+ is given by [18, Theorem 6.1].

Corollary 3.3 (Poincaré duality)**.** *A Schubert class ξ_w is Schur rigid if and only if its Poincaré dual ξ_{w^*} is Schur rigid.*

Proof. Given $w \in W^{\mathfrak{p}}$, let $w^* \in W^{\mathfrak{p}}$ be the element associated to the Poincaré dual of ξ_w. By [18, Corollary 6.2], the condition $\mathsf{H}_+(w)$ holds if and only if $\mathsf{H}_+(w^*)$ holds. It follows from Theorem 3.1 (resp. Corollary 3.2) that the Schubert system $\mathcal{B}_w$ (resp. the Schur system $\mathcal{R}_w$) is rigid if and only if the system $\mathcal{B}_{w^*}$ (resp. $\mathcal{R}_{w^*}$) is rigid. □

As given by [18, Definitions 5.27 and 5.36], we have

Definition 3.4. The condition H_1 *fails* when there exist $\mathfrak{g}_{0,0}$–highest weights $-\beta \in \Delta(\mathfrak{g}_{0,-1})$ and $\gamma \in \Delta(\mathfrak{g}_{1,\mathsf{a}})$ such that

$$[\mathfrak{g}_{-\beta}\,,\,\mathfrak{g}_{\gamma}] \overset{\text{(i)}}{=} \{0\} \neq [\mathfrak{g}_{\beta}\,,\,\mathfrak{g}_{\gamma}] \overset{\text{(ii)}}{=} [\mathfrak{g}_{\beta}\,,\,\mathfrak{g}_{1,\mathsf{a}}]\,. \tag{3.5}$$

The condition H_2 *fails* when there exist $\mathfrak{g}_{0,0}$–highest weights $\varepsilon \in \Delta(\mathfrak{g}_{1,\mathsf{a}-1})$ and $\gamma \in \Delta(\mathfrak{g}_{1,\mathsf{a}})$ such that

$$\{0\} \overset{\text{(i)}}{\neq} [\mathfrak{g}_{\varepsilon}\,,\,\mathfrak{g}_{-\gamma}] \overset{\text{(ii)}}{=} [\mathfrak{g}_{\varepsilon}\,,\,\mathfrak{g}_{-1,-\mathsf{a}}]\,. \tag{3.6}$$

When both H_1 and H_2 hold, we say that *the condition H_+ is satisfied.*

Remark. Note that $-\beta$ is a $\mathfrak{g}_{0,0}$–highest weight if and only if β is a $\mathfrak{g}_{0,0}$–lowest weight. That is, $\beta \in \Delta(\mathfrak{g}_{0,1})$ is a simple root.

Remark 3.7. Note that H_2 is trivially satisfied when $\mathsf{a} = 0$, as $\mathfrak{g}_{1,\mathsf{a}-1} = \mathfrak{g}_{1,-1} = 0$.

Remark 3.8. If the Dynkin diagram of G is simply laced (G is of type ADE), then H_2 is satisfied for every Schubert variety $X_w \subset X \simeq G/P_{\mathtt{i}}$. See Lemmas 4.4, 4.8, 4.19, and Remark 4.25(a).

Example. In Example 2.18, the $\mathfrak{g}_{0,0}$–lowest weights $\beta \in \Delta(\mathfrak{g}_{0,1})$ are $\{\alpha_2\,,\,\alpha_3\,,\,\alpha_7\,,\,\alpha_9\,,\,\alpha_{12}\}$, and the $\mathfrak{g}_{0,0}$–highest weights are

$$\begin{aligned} \Pi(\mathfrak{g}_{1,\mathsf{a}}) &= \{\alpha_1 + \cdots + \alpha_6\,,\ \alpha_3 + \cdots + \alpha_8\,,\ \alpha_4 + \cdots + \alpha_{11}\}\,,\\ \Pi(\mathfrak{g}_{1,\mathsf{a}-1}) &= \{\alpha_3 + \cdots + \alpha_6\,,\ \alpha_4 + \cdots + \alpha_8\}\,. \end{aligned}$$

The condition H_1 fails for $\beta = \alpha_9$ and $\gamma = \alpha_3 + \cdots + \alpha_8$. The condition H_2 is satisfied.

Example. In Example 2.27, the $\mathfrak{g}_{0,0}$–lowest weights $\beta \in \Delta(\mathfrak{g}_{0,1})$ are $\{\alpha_1\,,\,\alpha_2\,,\,\alpha_4\}$, and the $\mathfrak{g}_{0,0}$–highest weights are

$$\begin{aligned} \Pi(\mathfrak{g}_{1,\mathsf{a}}) &= \{\alpha_1 + \cdots + \alpha_5\,,\ \alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5\}\,,\\ \Pi(\mathfrak{g}_{1,\mathsf{a}-1}) &= \{\alpha_2 + \cdots + \alpha_5\,,\ 2\alpha_3 + 2\alpha_4 + \alpha_5\}\,. \end{aligned}$$

The condition H_1 fails for $\beta = \alpha_4$ and $\gamma = \alpha_1 + \cdots + \alpha_5$. The condition H_2 fails for $\varepsilon = 2\alpha_3 + 2\alpha_4 + \alpha_5$ and $\gamma = \varepsilon + \alpha_2$.

3.1. Outline of the proof of Theorem 3.1.

By [18, Theorem 5.38], the Schubert system $\mathcal{B}_w$ is rigid when H_+ is satisfied. So to prove Theorem 3.1 if suffices to show that there exist nontrivial integrals Y of the Schubert system when H_+ fails. This is done in Sections 3.2 and 3.4. We will see that there exists a Schubert divisor $X_{w'} \subset X_w$ and a 1-parameter subgroup $A \subset G$ with the property that $Y = \overline{A \cdot X_{w'}}$ is a nontrivial integral variety of the Schubert system $\mathcal{B}_w$.

To see that the varieties Y constructed in Sections 3.2 and 3.4 are nontrivial integrals of the Schubert system we must review some of the results of [18]. Define

$$\mathfrak{n}_w^\perp \ = \bigoplus_{\alpha \in \Delta(\mathfrak{g}_1) \backslash \Delta(w)} \mathfrak{g}_{-\alpha} \ = \ \bigoplus_{b > \mathsf{a}} \mathfrak{g}_{-1,-b} \,,$$

so that $\mathfrak{g}_{-1} = \mathfrak{n}_w \oplus \mathfrak{n}_w^\perp$, and let

$$\mathfrak{g}_w^\perp \ = \ \mathfrak{n}_w^\perp \ \oplus \ \mathfrak{g}_{0,-} \ \oplus \ \mathfrak{g}_{1,<\mathsf{a}} \,.$$

Each cohomology class $[\nu] \in H^1(\mathfrak{n}_w, \mathfrak{g}_w^\perp)$ admits a unique harmonic representative $\nu \in \mathcal{H}^1 \subset \mathfrak{g}_w^\perp \otimes \mathfrak{n}_w^*$ by [18, Proposition 5.10]. The grading element $Z_{\mathtt{i}}$ induces a graded decomposition $\mathcal{H}^1 = \mathcal{H}^1_0 \oplus \mathcal{H}^1_1 \oplus \mathcal{H}^1_2$ with $\mathcal{H}_0 \subset \mathfrak{n}_w^\perp \otimes \mathfrak{n}_w^*$, $\mathcal{H}^1_1 \subset \mathfrak{g}_{0,-} \otimes \mathfrak{n}_w^*$ and $\mathcal{H}^1_2 \subset \mathfrak{g}_{1,<\mathsf{a}} \otimes \mathfrak{n}_w^*$.

Let ϑ denote the $\mathfrak{g}$–valued, left-invariant Maurer-Cartan form on G. Set $\mathfrak{g}_w = \mathfrak{n}_w \oplus \mathfrak{g}_{0,\geq 0} \oplus \mathfrak{g}_{1,\geq \mathsf{a}}$, so that $\mathfrak{g} = \mathfrak{g}_w \oplus \mathfrak{g}_w^\perp$. Let $\vartheta_{k,\ell}$ denote the $\mathfrak{g}_{k,\ell}$–valued component of the Maurer-Cartan form, with respect to the direct sum decomposition $\mathfrak{g} = \oplus \mathfrak{g}_{k,\ell}$ of (2.13). Similarly, let $\vartheta_{\mathfrak{g}_w}$ and $\vartheta_{\mathfrak{n}_w^\perp}$ respectively denote the $\mathfrak{g}_w$– and $\mathfrak{n}_w^\perp$–valued components of ϑ, with respect to the direct sum decompositions $\mathfrak{g} = (\mathfrak{n}_w \oplus \mathfrak{n}_w^\perp) \oplus \mathfrak{g}_{\geq 0} = \mathfrak{g}_w \oplus \mathfrak{g}_w^\perp$. The following is an amalgam of [18, Lemma 4.9 and Corollary 5.12].

Lemma 3.9. *There is a bijective correspondence between submanifolds $U \subset G$ such that*

(i) *$\vartheta_{\mathfrak{g}_w} : T_g U \to \mathfrak{g}_w$ is a linear isomorphism for all $g \in U$,*
(ii) *$\vartheta_{\mathfrak{n}_w^\perp} = 0$ vanishes on U,*
(iii) *there exists a smooth map $\lambda : U \to \mathcal{H}^1_1$ such that $\vartheta_{0,-} = \lambda(\vartheta_{\mathfrak{n}_w})$ on U,*

and integral manifolds $U \cdot o \subset X$ of the Schubert system. Additionally, there exists a smooth function $\mu : U \to \mathfrak{g}_{1,<\mathsf{a}} \otimes \mathfrak{n}_w^*$ such that $\vartheta_{1,<\mathsf{a}} = \mu(\vartheta_{\mathfrak{n}_w})$ on U. If λ is identically zero, then μ takes values in $\mathcal{H}^1_2$.

The submanifold U of Lemma 3.9 is the sub-bundle $\mathcal{F}^0$ of [18, Corollary 5.12]. It is a consequence of the arguments of [18, Section 4.7] that the sub-bundle $\mathcal{F}^0$ is unique. Then [18, Proposition 4.13] yields

Lemma 3.10. *The integral manifold $U \cdot o \subset X$ is a Schubert variety of type w if and only if both λ and μ vanish identically on U.*

The integrals $Y = \overline{A \cdot X_{w'}}$ constructed in Sections 3.2 and 3.4 will be of the form $Y = \overline{U \cdot o}$, where U satisfies the hypothesis of Lemma 3.11.

Lemma 3.11. *Let $U \subset G$ be a submanifold such that*

(i) *$\vartheta_{\mathfrak{g}_w} : T_g U \to \mathfrak{n}_w$ is a linear isomorphism for all $g \in U$,*

(ii) $\vartheta_{\mathfrak{n}_w^\perp}$ *vanishes on* U.

Then $U \cdot o \subset X$ *is a* nontrivial *integral manifold of the Schubert system if either*

(a) *there exists a smooth map* $\lambda : U \to \mathcal{H}^1_1$, *that is not identically zero, such that* $\vartheta_{0,-} = \lambda(\vartheta_{\mathfrak{n}_w})$ *on* U; *or*

(b) $\vartheta_{0,-}$ *vanishes on* U, *and there exists a smooth map* $\mu : U \to \mathcal{H}^1_2$, *that is not identically zero, such that* $\vartheta_{1,<\mathsf{a}} = \mu(\vartheta_{\mathfrak{n}_w})$.

Proof. Let $G' \subset G$ be the connected Lie group with Lie algebra $\mathfrak{g}_{0,\geq 0} \oplus \mathfrak{g}_{1,\geq \mathsf{a}}$. Let $U \subset G$ be a submanifold satisfying (i) and (ii) of Lemma 3.11. Then $U' = UG' \subset G$ is a submanifold satisfying (i) and (ii) of Lemma 3.9.

Suppose U satisfies Lemma 3.11(a). Then U' satisfies Lemma 3.9(iii). Lemma 3.10 implies that $U \cdot o = U' \cdot o \subset X$ is a nontrivial integral of the Schubert system.

Suppose U satisfies Lemma 3.11(b). The $\vartheta_{0,-}$ also vanishes on U'. In particular, U' satisfies Lemma 3.9(iii) trivially. Moreover, since λ is identically zero on U', the function μ of Lemma 3.9 necessarily takes values in $\mathcal{H}^1_2$. Since μ is nonzero on $U \subset U'$, Lemma 3.10 implies that $U \cdot o = U' \cdot o \subset X$ is a nontrivial integral of the Schubert system. □

3.2. **Proof of Theorem 3.1 in the case that H_1 fails.** We will follow the strategy outlined in Section 3.1. Suppose that H_1 fails for a pair γ, $-\beta$; see Definition 3.4. Then [18, Lemma 5.28] asserts that

$$\mathfrak{g}_{-\beta} \otimes \mathfrak{g}_\gamma \subset \mathcal{H}^1_1 \,. \tag{3.12}$$

Lemma 2.41 and the fact that $\gamma \in \Delta(\mathfrak{g}_{1,\mathsf{a}})$ is a $\mathfrak{g}_{0,0}$–highest weight yield $w' \in W^{\mathfrak{p}}$ such that $\Delta(w') \sqcup \{\gamma\} = \Delta(w)$. Fix $0 \neq C \in \mathfrak{g}_{-\gamma}$ and $0 \neq B \in \mathfrak{g}_{-\beta}$. Let $\mathfrak{a} = \langle B + C \rangle$.

Claim. The direct sum $\mathfrak{n}' := \mathfrak{a} \oplus \mathfrak{n}_{w'}$ is a subalgebra of $\mathfrak{g}$.

The claim is proved at the end of this section. Assuming the claim holds, let $N' = \exp(\mathfrak{n}') \subset G$. When restricted to N' the Maurer-Cartan form ϑ takes values in $\mathfrak{n}'$. In particular, $U = N'$ satisfies hypotheses (i) and (ii) of Lemma 3.11. Moreover, the map λ of Lemma 3.11 is the nonzero, constant map given by $\lambda_{|\mathfrak{n}_{w'}} = 0$ and $\lambda(C) = B$. By (3.12), λ lies in $\mathcal{H}^1_1$. It follows from Lemma 3.11.a that $N' \cdot o \subset X$ is a nontrivial integral manifold of the Schubert system.

Finally, note that $\mathfrak{n}' \subset \mathfrak{g}_{-1} \oplus \mathfrak{g}_{0,-1}$ is nilpotent. It follows that $N' = \exp(\mathfrak{n}')$ is a linear algebraic subgroup of G. Therefore, the Zariski closure $Y_1 = \overline{N' \cdot o}$ is an irreducible algebraic subvariety of X with $\dim Y_1 = \dim N' = \dim X_w$. Therefore, Y_1 is a nontrivial, integral variety of the Schubert system. Modulo the claim, this completes the proof of Theorem 3.1 in the case that H_1 fails.

Remark. Let $A = \exp(\mathfrak{a})$. Then $\overline{N'} = \overline{A\, N_{w'}}$ and $Y_1 = \overline{A \cdot X_{w'}}$.

The proof the the claim makes use of the following

Remark 3.13. Given two roots $\mu, \nu \in \Delta$, we have $[\mathfrak{g}_\mu, \mathfrak{g}_\nu] = \mathfrak{g}_{\mu+\nu}$. In particular, $\mu + \nu \in \Delta$ if and only if $[\zeta, \xi] \neq 0$ for every $0 \neq \zeta \in \mathfrak{g}_\mu$ and $0 \neq \xi \in \mathfrak{g}_\nu$. See, for example, [13, Corollary 2.35].

Proof of claim. Since both $\mathfrak{a}$ and $\mathfrak{n}_{w'}$ are subalgebras of $\mathfrak{g}$, it suffices to show that $[\mathfrak{a}\,,\,\mathfrak{n}_{w'}] \subset \mathfrak{n}'$. To that end, let $\xi \in \mathfrak{n}_{w'}$. Then $C \in \mathfrak{g}_{-\gamma} \subset \mathfrak{g}_{-1}$ and Remark 2.3 yield

$$[B + C\,,\,\xi] \;=\; [B\,,\,\xi]\,.$$

Without loss of generality we may assume that $\xi \in \mathfrak{g}_{-\mu}$ is a nonzero root vector, $\mu \in \Delta(w')$. By Remark 3.13, $[B\,,\,\xi] \in \mathfrak{g}_{-\beta-\mu}$ is nonzero if and only if $\beta + \mu$ is a root; assume this is the case. We wish to show that $\mathfrak{g}_{-\beta-\mu} \subset \mathfrak{n}_{w'}$; equivalently, $\beta + \mu \in \Delta(w')$.

By Proposition 2.14, $\Delta(w) = \Delta(\mathfrak{g}_{1,\le \mathsf{a}})$. Since $\Delta(w') \sqcup \{\gamma\} = \Delta(w)$ and $\gamma \in \Delta(\mathfrak{g}_{1,\mathsf{a}})$, we have

$$0 \le \mu(Z_w) \le \mathsf{a} \quad \text{and} \quad \Delta(\mathfrak{g}_{1,<\mathsf{a}}) \;\subset\; \Delta(w') \;\subset\; \Delta(\mathfrak{g}_{1,\le\mathsf{a}})\,.$$

We consider three cases: First, if $\mu(Z_w) < \mathsf{a} - 1$, then $(\mu+\beta)(Z_w) < \mathsf{a}$ and $\beta + \mu \in \Delta(w')$. Second, if $\mu(Z_w) = \mathsf{a} - 1$, then $(\mu + \beta)(Z_w) = \mathsf{a}$. By (3.5.i), $\mu + \beta \neq \gamma$, so $\mu + \beta \in \Delta(w')$. Third, if $\mu(Z_w) = \mathsf{a}$, then (3.5.ii) implies $\beta + \mu$ is not a root. □

3.3. **The nontrivial Schubert integral Y_1.** Let Y_1 denote the nontrivial integral variety of the Schubert system $\mathcal{B}_w$ constructed in Section 3.2. By Corollary 2.9, the homology class $[Y_1]$ is an integer multiple of the Schubert class ξ_w. In Section 4 we give an algebro-geometric description of Y_1 and and will see that $[Y_1] = 2\xi_w$, in the case that X is one of the classical irreducible cominuscule varieties. The key tool in Section 4 is Lemma 3.17, which it is convenient to prove while the construction of Y_1 is fresh. We will obtain Proposition 3.15, which provides some coarse information about Y_1, as a corollary.

Let w, w', γ and β be as in Section 3.2. Let $(\cdot,\cdot)$ denote the Killing form on $\mathfrak{h}^*$, and define

$$\mathsf{s} \;:=\; 1 - 2\frac{(\gamma,\beta)}{(\beta,\beta)}\,. \tag{3.14}$$

Condition (3.5) and [13, Proposition 2.48] imply that the β–string containing γ is of the form $\gamma + \ell\beta$ with $0 \le \ell \le \mathsf{s} - 1$, and that $2 \le \mathsf{s} \le 4$. Given $0 \le j < \mathsf{s}$, set $\gamma_{j+1} := \gamma + j\beta$ and define

$$w^1 \;=\; r_{\gamma_{\mathsf{s}}} \cdots r_{\gamma_2} w \;\in\; W\,.$$

Proposition 3.15. *The Weyl group element w^1 is a member of the Hasse diagram $W^{\mathfrak{p}}$, and $\Delta(w^1) = \Delta(w) \sqcup \{\gamma_2, \ldots, \gamma_{\mathsf{s}}\}$. Let X_{w^1} denote the corresponding Schubert variety. Let Y_1 be the nontrivial integral of the Schubert system constructed in Section 3.2. Both Y_1 and X_w are subvarieties of X_{w^1} of codimension $\mathsf{s} - 1$.*

The proposition is proved below, after Example 3.19.

Let $C \in \mathfrak{g}_{-\gamma}$ and $B \in \mathfrak{g}_{-\beta}$ be as in Section 3.2. Set $C_1 = C$ and define $C_{\ell+1} = [B, C_\ell] \in \mathfrak{g}_{-\gamma-\ell\beta}$. Note that, $C_{\mathsf{s}} \neq 0$ and $C_{\mathsf{s}+1} = 0$. Consider the curve $\mathrm{C}_1 : \mathbb{C} \to \mathfrak{n}_{w'}^{\perp} \subset \mathfrak{g}_{-1}$ given by

$$\mathrm{C}_1(t) \;:=\; \sum_{\ell=1}^{\mathsf{s}} \frac{t^\ell}{\ell!} C_\ell\,.$$

Define

$$M_1 \;:=\; \exp(\mathfrak{m}_1) \quad \text{and} \quad \mathfrak{m}_1 \;:=\; \{\zeta + \mathrm{C}_1(t) \mid \zeta \in \mathfrak{n}_{w'}\,,\ t \in \mathbb{C}\}\,. \tag{3.16}$$

Lemma 3.17. *The nontrivial Schubert integral constructed in Section 3.2 is $Y_1 = \overline{M_1 \cdot o}$.*

The lemma is proved at the end of the section.

Remark 3.18. When $X = \mathrm{Gr}(k,n)$ and ξ_w is Schur flexible, Coskun has exhibited explicit irreducible varieties $Y \subset \mathrm{Gr}(k,n)$ such that $[Y] = 2\xi_w$ (cf. the proof of Theorem 2.1 in [7]). In the proof of Proposition 4.9, we will see that the nontrivial integrals Y_1 constructed in the proof above are of the form demonstrated by Coskun, as the following simple example illustrates.

Example 3.19. The notational conventions of Section 2.6 hold here. Let $X = \mathrm{Gr}(3,7) \subset \mathbb{P}(\bigwedge^3\mathbb{C}^7)$. The Schubert variety $X_w = \overline{\exp(\mathfrak{n}_w)\cdot o}$ given by $\mathsf{a} = 1$ and $\mathsf{J} = \{1,2,5\}$ has associated partition (cf. Section 2.6) $\lambda(w) = (1,4,7)$ is given by . The subspace $\mathfrak{n}_w = \{s^j_k\, e^k_j \mid s = (s^4_2, s^5_2, s^4_2, s^5_3, s^6_3, s^7_3) \in \mathbb{C}^6\}$ is represented in matrix form as

$$\mathfrak{n}_w \;=\; \begin{pmatrix} 0 & s^4_2 & s^4_3 \\ 0 & s^5_2 & s^5_3 \\ 0 & 0 & s^6_3 \\ 0 & 0 & s^7_3 \end{pmatrix} \;\subset\; \mathfrak{g}_{-1}\,,$$

We have $o = [e_1\wedge e_2\wedge e_3] \in \mathrm{Gr}(3,7)$. Thus $\exp(\mathfrak{n}_w)\cdot o = \{[e_1\wedge(e_2+s^j_2 e_j)\wedge(e_3+s^k_3 e_k)] \mid (s^j_k) \in \mathbb{C}^6\}$. Define $F^1 = \langle e_1\rangle$ and $F^4 = \langle e_1, e_2, e_4, e_5\rangle$. Then

$$X_w \;=\; \overline{\exp(\mathfrak{n}_w)\cdot o} \;=\; \left\{E \in \mathrm{Gr}(3,7) \mid F^1 \subset E\,,\ \dim(E\cap F^4) \ge 2\right\}\,.$$

The condition H_1 fails for the pair $\gamma = \alpha_2+\alpha_3+\alpha_4$ and $\beta = \alpha_5$. Set $C = e^2_5 \in \mathfrak{g}_{-\gamma}$ and $B = -e^5_6 \in \mathfrak{g}_{-\beta}$, then $C_2 = e^2_6$ and $C_3 = 0$. Thus, the integer of (3.14) is $\mathsf{s} = 2$, and $\mathrm{C}_1(t) = te^2_5 + \frac{1}{2}t^2 e^2_6$. So the space $\mathfrak{m}_1$ of (3.16) is represented in matrix form as

$$\mathfrak{m}_1 \;=\; \begin{pmatrix} 0 & s^4_2 & s^4_3 \\ 0 & t & s^5_3 \\ 0 & \frac{1}{2}t^2 & s^6_3 \\ 0 & 0 & s^7_3 \end{pmatrix} \;\subset\; \mathfrak{g}_{-1}\,.$$

Set $F^5 = \langle e_1, e_2, e_4, e_5, e_6\rangle$. The Schubert variety X_{w^1} of Proposition 3.15 is

$$X_{w^1} \;=\; \{E \in \mathrm{Gr}(3,7) \mid F^1 \subset E\,,\ \dim(E\cap F^5) \ge 2\}\,,$$

with associated partition $\lambda(w^1) = (1,5,7)$. Let $Q = \{(x_5)^2 - 2x_2x_6 = 0\} \subset F^5$. It follows from Lemma 3.17 that

$$Y_1 = \overline{\{E \in X_{w^1} \mid \exists\, \zeta \in \mathrm{Gr}(2,E) \text{ s.t. } \zeta \subset Q\}}\,.$$

This is the variety Y constructed by Coskun; he also shows that Y is irreducible and homologous to $2\xi_w$.

Proof of Proposition 3.15. Assume that Lemma 3.17 holds. Suppose that

$$w^1 \in W^{\mathfrak{p}} \quad\text{and}\quad \Delta(w^1) = \Delta(w) \sqcup \{\gamma_2, \ldots \gamma_{\mathsf{s}}\}\,. \tag{3.20}$$

Then it is immediate from $\Delta(w) \subset \Delta(w^1)$ and (2.12) that X_w is a codimension $\mathsf{s}-1$ subvariety of X_{w^1}. By (3.16), we have $\mathfrak{m}_1 \subset \mathfrak{n}_{w^1}$. Lemma 3.17 implies Y_1 is also a codimension $\mathsf{s}-1$ subvariety of X_{w^1}. So to prove the proposition it remains to establish (3.20). The argument is by induction.

Set $\Phi_1 = \Delta(w)$. Given $2 \le \ell \le \mathsf{s}$, set $\Phi_\ell = \Delta(w) \sqcup \{\gamma_2, \gamma_3, \ldots, \gamma_\ell\}$. Fix $\ell < \mathsf{s}$ and suppose that $\Phi_\ell = \Delta(\hat{w})$ for some $\hat{w} \in W^{\mathfrak{p}}$. (This inductive hypothesis holds for $\ell = 1$.) Let $\mathfrak{g} = \oplus \hat{\mathfrak{g}}_{j,k}$ be the $(Z_{\mathtt{i}}, Z_{\hat{w}})$–bigraded decomposition of $\mathfrak{g}$ given by (2.13).

Claim. The root $\gamma_{\ell+1}$ is a lowest $\hat{\mathfrak{g}}_{0,0}$–weight.

Given the claim, it follows from Lemma 2.42 that $r_{\gamma_{\ell+1}} \hat{w} \in W^{\mathfrak{p}}$ and $\Delta(r_{\gamma_{\ell+1}} \hat{w}) = \Phi_{\ell+1}$. This will complete the induction and the proof of Proposition 3.15.

To prove the claim, it suffices to show that

$$[\mathfrak{g}_{\gamma_{\ell+1}}, \mathfrak{g}_{-\alpha}] \ = \ 0 \tag{3.21}$$

for any simple root $\alpha \in \Delta(\hat{\mathfrak{g}}_{0,0})$.

Remark 3.13 and $\gamma_{\ell+1} = \gamma_\ell + \beta$ imply $\mathfrak{g}_{\gamma_{\ell+1}} = [\mathfrak{g}_{\gamma_\ell}, \mathfrak{g}_\beta]$. The Jacobi identity yields

$$[\mathfrak{g}_{\gamma_{\ell+1}}, \mathfrak{g}_{-\alpha}] \ = \ [\,[\mathfrak{g}_{\gamma_\ell}, \mathfrak{g}_\beta]\,,\, \mathfrak{g}_{-\alpha}] \ = \ [\,[\mathfrak{g}_{-\alpha}, \mathfrak{g}_\beta]\,,\, \mathfrak{g}_{\gamma_\ell}] \ + \ [\,[\mathfrak{g}_{\gamma_\ell}, \mathfrak{g}_{-\alpha}]\,,\, \mathfrak{g}_\beta]\ .$$

To see that $[\mathfrak{g}_{-\alpha}, \mathfrak{g}_\beta] = 0$, first note that $\beta + \gamma_\ell = \gamma_{\ell+1} \not\in \Phi_\ell = \Delta(\hat{w})$ implies that $\mathfrak{g}_\beta$ does not preserve $\mathfrak{n}_{\hat{w}}$. In particular, β is not a root of $\hat{\mathfrak{g}}_{0,0}$; so, $\alpha \neq \beta$. Since both α and β are simple, $-\alpha + \beta$ cannot a root; therefore, $[\mathfrak{g}_{-\alpha}, \mathfrak{g}_\beta] = 0$. Thus,

$$[\mathfrak{g}_{\gamma_{\ell+1}}, \mathfrak{g}_{-\alpha}] \ = \ [\,[\mathfrak{g}_{\gamma_\ell}, \mathfrak{g}_{-\alpha}]\,,\, \mathfrak{g}_\beta]\ .$$

Whence, $[\mathfrak{g}_{\gamma_{\ell+1}}, \mathfrak{g}_{-\alpha}] \neq 0$ only if $[\mathfrak{g}_{\gamma_\ell}, \mathfrak{g}_{-\alpha}] \neq 0$. As above, $[\mathfrak{g}_{\gamma_\ell}, \mathfrak{g}_{-\alpha}] = [[\mathfrak{g}_{\gamma_{\ell-1}}, \mathfrak{g}_{-\alpha}]\,,\, \mathfrak{g}_\beta]$. Continuing we see that

$$[\mathfrak{g}_{\gamma_{\ell+1}}, \mathfrak{g}_{-\alpha}] \neq 0 \quad \text{only if} \quad [\mathfrak{g}_{\gamma_{k+1}}, \mathfrak{g}_{-\alpha}] = [[\mathfrak{g}_{\gamma_k}, \mathfrak{g}_{-\alpha}], \mathfrak{g}_\beta] \neq 0 \tag{3.22}$$

for all $1 \le k \le \ell$.

Since α is a simple root, either $\alpha(Z_w) = 0$ or $\alpha(Z_w) = 1$. If $\alpha(Z_w) = 0$, then $[\mathfrak{g}_\gamma, \mathfrak{g}_{-\alpha}] \subset \mathfrak{g}_{1,\mathsf{a}}$. It follows from (3.5.ii) that $[[\mathfrak{g}_\gamma, \mathfrak{g}_{-\alpha}]\,,\, \mathfrak{g}_\beta] = 0$. Setting $k = 1$ in (3.22) yields (3.21). If $\alpha(Z_w) = 1$, then $[\mathfrak{g}_{\gamma_2}, \mathfrak{g}_{-\alpha}] = [[\mathfrak{g}_\gamma, \mathfrak{g}_{-\alpha}], \mathfrak{g}_\beta] \in \mathfrak{g}_{1,\mathsf{a}}$. Again, (3.5.ii) implies $[[\mathfrak{g}_{\gamma_2}, \mathfrak{g}_{-\alpha}]\,,\, \mathfrak{g}_\beta] = 0$, and setting $k = 2$ in (3.22) yields (3.21). Having established (3.21), the claim follows. □

Proof of Lemma 3.17. By definition $Y_1 = \overline{N' \cdot o}$, where $N' = \exp(\mathfrak{n}')$ and $\mathfrak{n}' = \mathfrak{n}_{w'} \oplus \langle C + B \rangle$. From the definition (3.16) of M_1, we see that to prove the lemma it suffices to show that

$$\exp t(C+B) \cdot \mathsf{v}_0 \ = \ (\exp \mathrm{C}_1(t)) \cdot \mathsf{v}_0\,, \quad \forall\ t \in \mathbb{C}\,. \tag{3.23}$$

Here $\mathsf{v}_0 \in V$ spans the highest $\mathfrak{g}$–weight line (so that $o = [\mathsf{v}_0]$).

Define $f(t) = \exp t(C+B) \cdot \mathsf{v}_0$ and $g(t) = \exp \mathrm{C}_1(t) \cdot \mathsf{v}_0$. To establish (3.23), it suffices to show that

$$f^{(k)}(0) \ = \ g^{(k)}(0) \tag{3.24}$$

for all $k \ge 0$. This may be done with a straight-forward induction argument as follows. First note that

$$f^{(k)}(0) \ = \ (C+B)^k \cdot \mathsf{v}_0\,.$$

Also

$$g^{(k)}(t) \ = \ \sum_{j=0}^{k-1} \left\{ \tbinom{k-1}{j} \left(\sum_{\ell=0}^{\mathsf{s}+j-k} \frac{t^\ell}{\ell!}\, C_{\ell+k-j} \right) g^{(j)}(t) \right\}\,,$$

so that

$$g^{(k)}(0) \;=\; \sum_{j=0}^{k-1} \tbinom{k-1}{j} C_{k-j}\, g^{(j)}(0)\,.$$

It is an exercise for the reader to verify that $(C+B)g^{(k)}(0) = g^{(k+1)}(0)$.

To complete the induction: (i) verify that (3.24) holds for $k = 0, 1$, and (ii) assume that (3.24) holds for $k \le n$, and compute $f^{(n+1)}(0) = (B+C)f^{(n)}(0) = (B+C)\,g^{(n)}(0) = g^{(n+1)}(0)$. □

3.4. **Proof of Theorem 3.1 in the case that H_2 fails.** We will follow the strategy outlined in Section 3.1. Suppose that H_2 fails for a pair $\gamma \in \Delta(\mathfrak{g}_{1,\mathsf{a}})$ and $\varepsilon \in \Delta(\mathfrak{g}_{1,\mathsf{a}-1})$, see Definition 3.4. Then [18, Lemma 5.39] asserts that

$$\mathfrak{g}_\varepsilon \otimes \mathfrak{g}_\gamma \;\subset\; \mathcal{H}^1_2\,. \tag{3.25}$$

By Lemma 2.41 there exists $w' \in W^{\mathfrak{p}}$ such that $\Delta(w') \sqcup \{\gamma\} = \Delta(w)$. Fix $0 \neq C \in \mathfrak{g}_{-\gamma}$ and $0 \neq E \in \mathfrak{g}_\varepsilon$. Set

$$\mathfrak{a} \;:=\; \langle C + E \rangle \;\subset\; \mathfrak{g} \qquad \text{and} \qquad A \;:=\; \exp(\mathfrak{a}) \;\subset\; G\,. \tag{3.26}$$

We will show that $Y_2 = \overline{AN_{w'} \cdot o}$ is a nontrivial integral variety of the Schubert system.

Because $\exp : \mathfrak{g} \to G$ is a diffeomorphism from a neighborhood of $0 \in \mathfrak{g}$ to a neighborhood of $\mathrm{Id} \in G$, it follows that there exist connected neighborhoods $A_0 \subset A$ and $N_0 \subset N_{w'}$ of Id that are embedded submanifolds in G. Let $\mathbf{m} : G \times G \to G$ denote the multiplication map. Then $\mathbf{m}_* : \mathfrak{g} \times \mathfrak{g} \to \mathfrak{g}$ is given by $\mathbf{m}_*(u,v) = u + v$. In particular, the restriction $\mathbf{m}_* : \mathfrak{a} \times \mathfrak{n}_{w'} \to \mathfrak{a} \oplus \mathfrak{n}_{w'}$ is a bijection. It follows, shrinking A_0 and N_0 if necessary, that there is a neighborhood $U \subset G$ of Id such that $\mathbf{m} : A_0 \times N_0 \to U$ is an isomorphism.

Compute the Maurer-Cartan form ϑ on $U \subset G$ as follows. Let $g_0 g_1 \in U$ with $g_0 \in A_0$ and $g_1 \in N_0$. The isomorphism $\phi : G \to G$ defined by $\phi(g) = g_0 g g_1$ preserves $A\,N_{w'}$ and so induces an isomorphism $\phi_* : \mathfrak{a} \oplus \mathfrak{n}_{w'} \to T_{g_0 g_1} U$. So any element of $T_{g_0 g_1}U$ may be expressed as $\phi_* \xi$ with $\xi \in \mathfrak{a} \oplus \mathfrak{n}_{w'}$. Given $g \in G$, let $L_g, R_g : G \to G$ denote the left- and right-multiplication maps. Then $\phi = L_{g_0}\, R_{g_1}$. By definition

$$\vartheta(\phi_* \xi) \;=\; (L^{-1}_{g_0 g_1})_*(\phi_* \xi) \;=\; \mathrm{Ad}_{g_1^{-1}} \xi\,. \tag{3.27a}$$

Therefore, $\vartheta_{|U}$ takes values in $\mathrm{Ad}_{N_{w'}}(\mathfrak{a} \oplus \mathfrak{n}_{w'}) = \mathrm{Ad}_{N_{w'}}(\mathfrak{a}) \oplus \mathrm{Ad}_{N_{w'}}(\mathfrak{n}_{w'})$. Since $\mathfrak{n}_{w'}$ is an algebra, we have

$$\mathrm{Ad}_{N_{w'}}(\mathfrak{n}_{w'}) \;=\; \mathfrak{n}_{w'}\,. \tag{3.27b}$$

It remains to consider

$$\mathrm{Ad}_{N_{w'}}(\mathfrak{a}) \;=\; \mathrm{Ad}_{\exp(\mathfrak{n}_{w'})}(\mathfrak{a}) \;=\; (\exp \circ \mathrm{ad}_{\mathfrak{n}_{w'}})(\mathfrak{a})\,. \tag{3.27c}$$

The algebra $\mathfrak{a}$ is spanned by $C+E$. By Remark 2.3, $[\mathfrak{n}_{w'}, C+E] = [\mathfrak{n}_{w'}, E]$. Equations (2.15, 3.6), and $\mathfrak{n}_{w'} \oplus \mathfrak{g}_{-\gamma} = \mathfrak{n}_w$ imply that $[\mathfrak{n}_{w'}\,,\, E] \;\subset\; \mathfrak{g}_0$. Moreover, the fact that $\varepsilon \in \Delta(\mathfrak{g}_{1,1-\mathsf{a}})$ is a highest $\mathfrak{g}_{0,0}$–weight yields

$$[\mathfrak{n}_{w'}\,,\, E] \;\subset\; \mathfrak{h}_\varepsilon \oplus \mathfrak{g}_0^+\,, \quad \text{where } \ \mathfrak{h}_\varepsilon := [\mathfrak{g}_\varepsilon\,,\, \mathfrak{g}_{-\varepsilon}] \ \text{ and } \ \mathfrak{g}_0^+ := \oplus_{\alpha \in \Delta^+(\mathfrak{g}_0)} \mathfrak{g}_\alpha\,.$$

By Proposition 2.14, $\Delta(w') \sqcup \{\gamma\} = \Delta(w) = \Delta(\mathfrak{g}_{1,\le \mathsf{a}})$. So, since $\gamma \in \Delta(\mathfrak{g}_{1,\mathsf{a}})$ is a highest $\mathfrak{g}_{0,0}$–weight, we have $[\mathfrak{n}_{w'}, \mathfrak{h}_\varepsilon \oplus \mathfrak{g}_0^+] \subset \mathfrak{n}_{w'}$. Therefore,

$$(\exp \circ \mathrm{ad}_{\mathfrak{n}_{w'}})(\mathfrak{a}) \ \subset\ \mathfrak{a} \ \oplus\ (\mathfrak{h}_\varepsilon \oplus \mathfrak{g}_0^+) \ \oplus\ \mathfrak{n}_{w'} \,. \tag{3.27d}$$

Now (3.27) yields $\vartheta_{\mathfrak{n}_w^\perp} = 0$ on U, and $\vartheta_{\mathfrak{n}_w} : T_zU \to \mathfrak{n}_w$ is a linear isomorphism for all $z \in U$. Lemma 3.11 implies $U \cdot o \subset X$ is an integral manifold of the Schubert system. Equations (3.27) also imply that $\vartheta_{0,-}$ vanishes on U. Similarly, (3.27) implies that $\vartheta_{1,<\mathsf{a}} = \mu(\vartheta_{\mathfrak{n}_w})$, where $\mu : U \to \mathfrak{g}_{1,<\mathsf{a}} \otimes \mathfrak{n}_w^*$ is the nonzero, constant map defined by $\mu_{|\mathfrak{n}_{w'}} = 0$ and $\mu(C) = E$. By (3.25), μ takes values in $\mathcal{H}^1_2$. By Lemma 3.11(b), $U \cdot o \subset X$ is a nontrivial integral manifold of the Schubert system.

Finally, to see that the Zariski closure

$$Y_2 \ =\ \overline{U \cdot o} \ =\ \overline{AN_{w'} \cdot o} \tag{3.28}$$

is a variety of dimension $|w|$, it suffices to observe that A and $N_{w'}$ are algebraic. This is a consequence of the fact that both $\mathfrak{a}$ and $\mathfrak{n}_{w'}$ are nilpotent. The nilpotency of $\mathfrak{n}_{w'}$ is immediate from Remark 2.3. To see that $\mathfrak{a}$ is nilpotent, note that $[\mathfrak{a}, \mathfrak{g}_{\pm 1}] \subset \mathfrak{g}_0$ and $\mathrm{ad}^2_{\mathfrak{a}}(\mathfrak{g}_{0,\ell}) \subset \mathfrak{g}_{0,\ell-1}$; these relations imply that the adjoint action of $\mathfrak{a}$ on $\mathfrak{g}$ is nilpotent. This completes the proof of Theorem 3.1.

3.5. The nontrivial Schubert integral Y_2.

Let Y_2 denote the nontrivial integral variety of the Schubert system $\mathcal{B}_w$ constructed in Section 3.4. By Corollary 2.9, the homology class $[Y_2]$ is an integer multiple of the Schubert class ξ_w. In Section 4 we give an algebro-geometric description of Y_2 and determine that homology multiple. The key tool is Lemma 3.36, which it is convenient to prove while the construction of Y_2 is fresh. We obtain Proposition 3.31, which provides some coarse information about Y_2, as a corollary of the lemma.

Let $\gamma \in \Delta(\mathfrak{g}_{1,\mathsf{a}})$, $\varepsilon \in \Delta(\mathfrak{g}_{1,\mathsf{a}-1})$ and $w, w' \in W^{\mathfrak{p}}$ be as in Section 3.4. Set

$$\beta \ :=\ \gamma - \varepsilon \ \in\ \Delta(\mathfrak{g}_{0,1}) \,.$$

Claim 3.29. Let $\gamma_2 = \gamma + \beta$. Then γ_2 is a root.

Claim 3.30. Let $w^2 = r_{\gamma_2} w \in W$. Then $w^2 \in W^{\mathfrak{p}}$ and $\Delta(w^2) = \{\gamma_2\} \sqcup \Delta(w)$.

Proposition 3.31. *Let X_{w^2} denote the Schubert variety associated with w^2, see* (2.12). *Given $t \in \mathbb{C}$ there exist varieties $Y_2(t) \subset X_{w^2}$ of dimension $|w'| = \dim X_w - 1$ such that $Y_2(0) = X_{w'}$ and*

$$Y_2 \ =\ \overline{\bigcup_{t \in \mathbb{C}} Y_2(t)} \ \subset\ X_{w^2} \,.$$

The proposition is proved below.

Proof of Claim 3.29. By Remark 3.13, γ_2 is a root if and only if $\mathfrak{g}_{\gamma_2} = [\mathfrak{g}_\gamma, \mathfrak{g}_\beta]$ is nonzero. By (3.6.ii), the Jacobi identity and Remark 2.3,

$$[\mathfrak{g}_\gamma, \mathfrak{g}_\beta] \ =\ [\mathfrak{g}_\gamma \,,\, [\mathfrak{g}_{1,\mathsf{a}} \,,\, \mathfrak{g}_{-\varepsilon}]] \ =\ [\mathfrak{g}_{1,\mathsf{a}} \,,\, [\mathfrak{g}_\gamma, \mathfrak{g}_{-\varepsilon}]] \ =\ [\mathfrak{g}_{1,\mathsf{a}} \,,\, \mathfrak{g}_\beta] \ .$$

So γ_2 fails to be a root if and only if $[\mathfrak{g}_{1,\mathsf{a}} \,,\, \mathfrak{g}_\beta] = 0$. If this is the case, then $-\beta \in \Delta(\mathfrak{g}_{0,-1})$ and Proposition 2.14 imply that $\mathfrak{g}_{-\beta}$ stabilizes $\mathfrak{n}_w = \mathfrak{g}_{-1,\ge -\mathsf{a}}$. But this contradicts the fact that the Lie subalgebra of $\mathfrak{g}_0$ stabilizing $\mathfrak{n}_w$ is $\mathfrak{g}_{0,\ge 0}$, see Section 2.4. So γ_2 must be a root. □

Proof of Claim 3.30. It is immediate from $\gamma \in \Delta(\mathfrak{g}_{1,\mathsf{a}})$ and $\beta \in \Delta(\mathfrak{g}_{0,1})$ that $\gamma_2 \in \Delta(\mathfrak{g}_{1,\mathsf{a}+1})$. So, by Lemma 2.42, to prove the claim, it suffices to show that γ_2 is a lowest $\mathfrak{g}_{0,0}$–weight. Equivalently, if $\alpha \in \Delta^+(\mathfrak{g}_{0,0})$ is a simple root, then $[\mathfrak{g}_{\gamma_2}, \mathfrak{g}_{-\alpha}] = 0$. By Remark 3.13 and the Jacobi identity, we have

$$[\mathfrak{g}_{\gamma_2}, \mathfrak{g}_{-\alpha}] \;=\; [[\mathfrak{g}_\gamma, \mathfrak{g}_\beta]\,,\, \mathfrak{g}_{-\alpha}] \;=\; [[\mathfrak{g}_{-\alpha}, \mathfrak{g}_\beta]\,,\, \mathfrak{g}_\gamma] + [[\mathfrak{g}_\gamma, \mathfrak{g}_{-\alpha}]\,,\, \mathfrak{g}_\beta]\,. \tag{3.32}$$

To see that $[\mathfrak{g}_{-\alpha}, \mathfrak{g}_\beta]$ is zero, note that Remark 3.13 and the Jacobi identity yield

$$[\mathfrak{g}_{-\alpha}, \mathfrak{g}_\beta] \;=\; [\mathfrak{g}_{-\alpha}\,,\, [\mathfrak{g}_\gamma, \mathfrak{g}_{-\varepsilon}]] \;=\; [\mathfrak{g}_\gamma\,,\, [\mathfrak{g}_{-\alpha}, \mathfrak{g}_{-\varepsilon}]] + [\mathfrak{g}_{-\varepsilon}\,,\, [\mathfrak{g}_\gamma, \mathfrak{g}_{-\alpha}]]\,.$$

The hypothesis that ε is a highest $\mathfrak{g}_{0,0}$–weight implies $[\mathfrak{g}_{-\varepsilon}, \mathfrak{g}_{-\alpha}] = 0$. The hypothesis that H_2 fails for the pair γ,ε implies $[\mathfrak{g}_{-\varepsilon}\,,\, [\mathfrak{g}_\gamma, \mathfrak{g}_{-\alpha}]] = 0$. Thus, $[\mathfrak{g}_{-\alpha}, \mathfrak{g}_\beta] = 0$. Therefore,

$$\begin{aligned}
[\mathfrak{g}_{\gamma_2}, \mathfrak{g}_{-\alpha}] \;&\overset{(3.32)}{=}\; [[\mathfrak{g}_\gamma, \mathfrak{g}_{-\alpha}]\,,\, \mathfrak{g}_\beta] \overset{(1)}{=} [\mathfrak{g}_{\gamma-\alpha}\,,\, [\mathfrak{g}_\gamma, \mathfrak{g}_{-\varepsilon}]]\\
&\overset{(2)}{=}\; [\mathfrak{g}_\gamma\,,\, [\mathfrak{g}_{\gamma-\alpha}, \mathfrak{g}_{-\varepsilon}]] + [\mathfrak{g}_{-\varepsilon}\,,\, [\mathfrak{g}_\gamma, \mathfrak{g}_{\gamma-\alpha}]]\,;
\end{aligned}$$

above, the equality (1) is a consequence of Remark 3.13 and $\beta = \gamma - \varepsilon$, and the equality (2) is the Jacobi identity. If $\gamma - \alpha$ is not a root, then the equation above and Remark 3.13 yield $[\mathfrak{g}_{\gamma_2}, \mathfrak{g}_{-\alpha}] = 0$. If $\gamma - \alpha$ is a root, then γ and $\gamma - \alpha$ are both roots of $\mathfrak{g}_{1,\mathsf{a}} \subset \mathfrak{g}_1$; so Remark 2.3 and the equation above yield $[\mathfrak{g}_{\gamma_2}, \mathfrak{g}_{-\alpha}] = [\mathfrak{g}_\gamma\,,\, [\mathfrak{g}_{\gamma-\alpha}, \mathfrak{g}_{-\varepsilon}]]$. Finally, the hypothesis that H_2 fails for the pair γ,ε yields $[\mathfrak{g}_{\gamma-\alpha}, \mathfrak{g}_{-\varepsilon}] = 0$. Whence $[\mathfrak{g}_{\gamma_2}, \mathfrak{g}_{-\alpha}] = 0$. □

With $E \in \mathfrak{g}_\varepsilon$ and $C \in \mathfrak{g}_{-\gamma}$ as in Section 3.4, set

$$\beta \;:=\; \gamma - \varepsilon \quad\text{and}\quad B \;:=\; [E, C] \;\in\; \mathfrak{g}_{-\beta} \;\subset\; \mathfrak{g}_{0,-1}\,. \tag{3.33}$$

Let

$$C_\flat \;:=\; [B, C] \;\in\; \mathfrak{g}_{-\gamma_2} \;\subset\; \mathfrak{g}_{-1}\,.$$

Set

$$\mathrm{C}_2(t) \;=\; tC - \tfrac{1}{3}t^3 C_\flat \quad\text{and}\quad \mathbf{b}(t) \;=\; \exp\left(\tfrac{1}{2}t^2 B\right)\,. \tag{3.34}$$

Define

$$M_2 \;:=\; \exp(\mathfrak{m}_2)\,, \quad\text{where}\quad \mathfrak{m}_2 \;:=\; \left\{\mathrm{Ad}_{\mathbf{b}(t)}\left(\mathrm{C}_2(t) + \zeta\right) \;|\; t \in \mathbb{C}\,,\; \zeta \in \mathfrak{n}_{w'}\right\}\,. \tag{3.35}$$

Lemma 3.36. *The nontrivial Schubert integral constructed in Section 3.4 is* $Y_2 = \overline{M_2 \cdot o}$.

Proof of Proposition 3.31. Given $t \in \mathbb{C}$, let

$$\mathfrak{m}_2(t) \;=\; \mathrm{Ad}_{\mathbf{b}(t)}\{\mathrm{C}_2(t) + \zeta \;|\; \zeta \in \mathfrak{n}_{w'}\} \tag{3.37}$$

and $M_2(t) = \exp\mathfrak{m}_2(t)$. Set

$$Y_2(t) \;=\; \overline{M_2(t) \cdot o}\,. \tag{3.38}$$

Then $Y_2(0) = X_{w'}$, and Lemma 3.36 yields $Y_2 = \overline{\cup_{t\in\mathbb{C}} Y_2(t)}$. □

Proof of Lemma 3.36. Set $E_\flat = [E, B] \in \mathfrak{g}_{\varepsilon-\beta} \subset \mathfrak{g}_1$. By Remark 2.3,

$$[C, C_\flat] \;=\; 0 \;=\; [E, E_\flat]\,. \tag{3.39a}$$

Since $\mathfrak{g}_{-2\beta} = \mathfrak{g}_{-3\beta} = \{0\}$, we also have

$$[C, E_\flat] \;=\; [E, C_\flat] \;=\; [C_\flat, E_\flat] \;=\; 0\,. \tag{3.39b}$$

I claim that

$$[B, C_\flat] \;=\; 0 \;=\; [E_\flat, B] \tag{3.39c}$$

as well. To see this, it is convenient to work in the universal enveloping algebra $U(\mathfrak{g})$. Let $C' = [B, C_\flat] \in \mathfrak{g}_{-\gamma-2\beta} \subset \mathfrak{g}_{-1}$. Then

$$\begin{aligned} &E\,C\,C_\flat \;\overset{(3.33)}{=}\; (C\,E \,+\, B)\,C_\flat \;\overset{(3.39b)}{=}\; C\,C_\flat\,E \,+\, C_\flat\,B \,+\, C' \qquad \text{and}\\ &E\,C\,C_\flat \;\overset{(3.39a)}{=}\; E\,C_\flat\,C \;\overset{(3.39b)}{=}\; C_\flat\,E\,C \;\overset{(3.33)}{=}\; C_\flat\,(C\,E \,+\, B) \;\overset{(3.39a)}{=}\; C\,C_\flat\,E \,+\, C_\flat\,B\,, \end{aligned}$$

from which it follows that $C' = 0$. A similar argument yields $[E_\flat, B] = 0$, establishing (3.39c).

The first equation of (3.39c) implies $\mathrm{C}(t) := \mathrm{Ad}_{\mathbf{b}(t)} C_2(t) = tC + \frac{1}{6}t^3 C_\flat$. So the expression for $\mathfrak{m}_2$ in (3.35) may be rewritten as

$$\mathfrak{m}_2 \;=\; \{\mathrm{C}(t) + \mathrm{Ad}_{\mathbf{b}(t)}\zeta \mid t \in \mathbb{C}\,,\ \zeta \in \mathfrak{n}_{w'}\}\,. \tag{3.40}$$

Let $\mathbf{c}(t) = \exp \mathrm{C}(t)$. Similarly, define $\mathrm{E}(t) = tE + \frac{1}{6}t^3 E_\flat$ and $\mathbf{e}(t) = \exp \mathrm{E}(t)$.

Claim 3.41. The one-parameter subgroup of (3.26) is $A = \{\mathbf{c}(t)\,\mathbf{b}(t)\,\mathbf{e}(t) \mid t \in \mathbb{C}\}$.

Assume the claim holds. Using $\mathrm{E}(t) \in \mathfrak{g}_1$, we have $\mathbf{e}(t)\,N_{w'} \cdot o = \mathbf{e}(t)\,N_{w'}\,\mathbf{e}(t)^{-1} \cdot o = N_{w'} \cdot o$. Similarly, since $B \in \mathfrak{g}_0$, we have

$$A\,N_{w'} \cdot o \;=\; \mathbf{c}(t)\,\mathbf{b}(t)\,N_{w'} \cdot o \;=\; \mathbf{c}(t)\,\mathbf{b}(t)\,N_{w'}\,\mathbf{b}(t)^{-1} \cdot o\,.$$

Now (3.35) and (3.40) yield $\overline{A\,N_{w'} \cdot o} = \overline{M_2 \cdot o}$. Finally, (3.28) yields Lemma 3.36.

It remains to prove Claim 3.41. Set $f(t) = \exp t(C+E)$ and $g(t) = \mathbf{c}(t)\,\mathbf{b}(t)\,\mathbf{e}(t)$. Since $A = \exp\{t(C+E) \mid t \in \mathbb{C}\}$, to prove the claim it suffices to show that $f(t) = g(t)$ for all $t \in \mathbb{C}$. Equivalently, $f^{(n)}(0) = g^{(n)}(0)$ for all n. Note that $f^{(n)}(0) \;=\; (C+E)^n$. So, it remains to show that $g^{(n)}(0) = (C+E)^n$.

Compute

$$\mathbf{b}(t) \;=\; \sum_{n=0}^{\infty} \frac{t^{2n}\,B^n}{2^n\,n!}\,, \quad \mathbf{c}(t) \;=\; \sum_{\substack{n=0\\ j+3k=n}}^{\infty} \frac{t^n\,C^j\,C_\flat^k}{6^k\,j!\,k!}\,, \quad \mathbf{e}(t) \;=\; \sum_{\substack{n=0\\ j+3k=n}}^{\infty} \frac{t^n\,E^j\,E_\flat^k}{6^k\,j!\,k!}\,,$$

so that

$$\begin{aligned} g^{(n)}(0) \;&=\; n! \sum_{j+k+2m=n} \frac{\mathbf{c}^{(j)}(0)\,\mathbf{b}^{(2m)}(0)\,\mathbf{e}^{(k)}(0)}{j!\,k!\,(2m)!}\\ &=\; \sum_{j+k+2m=n}\ \sum_{\substack{p_1+3p_2=j\\ q_1+3q_2=k}} \frac{C^{p_1}\,C_\flat^{p_2}\,B^m\,E^{q_1}\,E_\flat^{q_2}}{2^m\,6^{p_2+q_2}\,p_1!\,p_2!\,m!\,q_1!\,q_2!} \end{aligned}$$

For the computations that follow it is helpful to keep (3.39) in mind. The formulas

$$\begin{aligned} E\,B^p \;&=\; B^p\,E \,+\, p\,B^{p-1}\,E_\flat\,,\\ E\,C^p \;&=\; C^p\,E \,+\, p\,C^{p-1}\,B \,+\, \tbinom{p}{2}\,C^{p-2}\,C_\flat \end{aligned}$$

may be established by induction, and yield, after a tedious but straight-forward computation, $(C+E)\,g^{(n)}(0) = g^{(n+1)}(0)$. Inductively, this gives us $g^{(n)}(0) = (C+E)^n g(0) = (C+E)^n$. $\square$

Lemma 3.42. *The one-parameter subgroup* $\mathcal{B} = \{\exp tB \mid t \in \mathbb{C}\} \subset G$ *preserves the Schubert variety* X_{w^2} *of Proposition 3.31. In particular,* $\mathbf{b}(t) \in \mathcal{B}$*, defined by* (3.34)*, preserves* X_{w^2} *for all* $t \in \mathbb{C}$.

Proof. Let $b \in \mathcal{B}$. By definition (2.12), $X_{w^2} = \overline{N_{w^2} \cdot o}$. Since $B \in \mathfrak{g}_0$, we have $bN_{w^2} \cdot o = bN_{w^2}b^{-1} \cdot o$. Since $N_{w^2} = \exp(\mathfrak{n}_{w^2})$, we see that it suffices to show that Ad_b preserves $\mathfrak{n}_{w^2}$; equivalently, ad_B preserves $\mathfrak{n}_{w^2}$. Since $B \in \mathfrak{g}_{-\beta}$, this is equivalent to $[\mathfrak{g}_{-\beta}, \mathfrak{n}_{w^2}] \subset \mathfrak{n}_{w^2}$. From the definition (2.4) of $\mathfrak{n}_{w^2}$ and Remark 3.13, we see that ad_B preserves $\mathfrak{n}_{w^2}$ if and only if the following holds:

$$\text{If } \mu \in \Delta(w^2) \text{ and } \mu + \beta \text{ is a root, then } \mu + \beta \in \Delta(w^2).$$

By Claim 3.30, $\Delta(w^2) = \{\gamma_2\} \sqcup \Delta(w)$. If $\mu = \gamma_2$, then (3.39c) implies that $\mu + \beta$ is not a root. So we may suppose that $\mu \in \Delta(w)$. By Proposition 2.14, $\Delta(w) = \Delta(\mathfrak{g}_{1,\le \mathsf{a}})$. Recall that $\beta \in \Delta(\mathfrak{g}_{0,1})$, see (3.33). So, if $\mu \in \Delta(\mathfrak{g}_{1,<\mathsf{a}}) \subset \Delta(w)$ and $\mu + \beta$ is a root, then $\mu + \beta \in \Delta(w)$. It remains to consider the case that $\mu \in \Delta(\mathfrak{g}_{1,\mathsf{a}})$. If $\mu = \gamma$, then $\mu + \beta = \gamma_2 \in \Delta(w^2)$. So suppose that $\mu \in \Delta(\mathfrak{g}_{1,\mathsf{a}}) \backslash \{\gamma\}$. We wish to see that $\mu + \beta$ is not a root. Making use of Remarks 2.3 and 3.13 and the Jacobi identity, we have $[\mathfrak{g}_\mu, \mathfrak{g}_\beta] = [\mathfrak{g}_\mu, [\mathfrak{g}_\gamma, \mathfrak{g}_{-\varepsilon}]] = [\mathfrak{g}_\gamma, [\mathfrak{g}_\mu, \mathfrak{g}_{-\varepsilon}]]$. Since H_2 fails for the pair γ,ε, it must be the case that $[\mathfrak{g}_\mu, \mathfrak{g}_{-\varepsilon}] = 0$. Thus $[\mathfrak{g}_\mu, \mathfrak{g}_\beta] = 0$, and $\mu + \beta$ is not a root by Remark 3.13. □

4. The Schubert integrals

Let Y be one of the Schubert integrals Y_1 or Y_2 constructed in Sections 3.2 and 3.4. By Corollary 2.9, there exists an integer $r > 0$ such that $[Y] = r\xi_w$, where $\xi_w = [X_w]$ is the homology class represented by the Schubert variety X_w. In this section, we will give algebro-geometric descriptions of the integrals Y, and use the descriptions to determine the homology multiple r, in the case that X is an irreducible, classical cominuscule variety. For example, in many cases the integral Y_1 is the set of points in the Schubert variety X_{w^1} (of Proposition 3.15) satisfying a quadric condition. The main results of this section are Propositions 4.2, 4.5, 4.9, 4.13, 4.20. The key tools are Lemmas 3.17 and 3.36.

We will see that if $Y = Y_1$, then $r = 2$; and if $Y = Y_2$, then $r \in \{3, 4\}$. The integer r will be determined as follows. Let $w^* \in W^{\mathfrak{p}}$ be the dual of w, see [18, Section 2.5]. Then ξ_{w^*} is the Poincaré dual of ξ_w. By [12], the integer r is the number of points in the intersection $\widetilde{X}_{w^*} \cap Y$, where $\widetilde{X}_{w^*}$ is a general Schubert variety of type w^*.

4.1. Quadric hypersurfaces $\mathcal{Q}^m \subset \mathbb{P}^{m+1}$.

From [18, Theorem 6.1] and Theorem 3.1 we have the following.

Proposition 4.1. *Every Schubert class in an odd dimensional quadric* $\mathcal{Q}^{2n-1} = B_n/P_1$ *is Schubert flexible. The only Schubert rigid classes in an even dimensional quadric* $\mathcal{Q}^{2n-2} = D_n/P_1$ *are the two represented by maximal linear linear spaces* $\mathbb{P}^{n-1} \subset \mathcal{Q}^{2n-2}$*. They correspond to* $\mathsf{a} = 0$*,* $\mathsf{J} = \{n-1\}$ *and* $\mathsf{a} = 0$*,* $\mathsf{J} = \{n\}$.

4.1.1. *Odd dimensional quadrics* $\mathcal{Q}^{2n-1} = B_n/P_1$. Set $m = 2n - 1$. Review the notation and definitions of Section 2.5.1, and recall that $\mathsf{J} = \{\mathsf{j}\} \subset \{2, \ldots, n\}$ and $\mathsf{a} = 0, 1$. There are three cases to consider, see [18, Section 6.1].

(A) If $\mathsf{a} = 0$, then H_1 fails for $\gamma = \alpha_1 + \cdots + \alpha_{\mathsf{j}-1}$ and $\beta = \alpha_{\mathsf{j}}$. By Remark 3.7, the condition H_2 is trivial.

(B) If $\mathsf{a} = 1$ and $\mathsf{j} < n$, then H_1 fails for $\gamma = \alpha_1 + \cdots + \alpha_{\mathsf{j}} + 2(\alpha_{\mathsf{j}+1} + \cdots \alpha_n)$ and $\beta = \alpha_{\mathsf{j}}$ as long as $\mathsf{j} < n$. The condition H_2 is satisfied.[†]

(C) If $\mathsf{a} = 1$ and $\mathsf{j} = n$, then H_1 holds, but H_2 fails for $\gamma = \alpha_1 + \cdots + \alpha_n$ and $\varepsilon = \alpha_1 + \cdots + \alpha_{n-1}$.

Proposition 4.2. (a.i) *If $\mathsf{j} < n$, then nontrivial Schubert integral Y_1 constructed for Case (A) in Section 3.2 is a cone $Q \subset \mathbb{P}^{\mathsf{j}}$ over a conic with vertex $\mathbb{P}^{\mathsf{j}-3}$.*

(a.ii) *If $\mathsf{j} = n$, then there exist a Schubert variety $X_{w^1} = \mathcal{Q}^m \cap \mathbb{P}^{n+2}$ containing $X_w = \mathbb{P}^{n-1}$ as a codimension two subvariety, and cones $Q, R \subset \mathbb{P}^{n+2}$ over a conic and a cubic, such that the Schubert integral constructed for Case (A) in Section 3.2 is $Y_1 = X_{w^1} \cap Q \cap R$.*

(b) *There exist a Schubert variety $X_{w^1} = \mathcal{Q}^m \cap \mathbb{P}^{2n-\mathsf{j}+2}$ containing $X_w = \mathcal{Q}^m \cap \mathbb{P}^{2n-\mathsf{j}+1}$ as a codimension one subvariety, and a cone $Q \subset \mathbb{P}^{2n-\mathsf{j}+2}$ over a conic (with vertex $\mathbb{P}^{2n-\mathsf{j}-1}$) such that the Schubert integral constructed for Case (B) in Section 3.2 is $Y_1 = X_{w^1} \cap Q$.*

(c) *There exists a Schubert variety $X_{w^2} = \mathcal{Q}^m \cap \mathbb{P}^{n+2}$, containing $X_w = \mathcal{Q}^m \cap \mathbb{P}^{n+1}$ as a codimension one subvariety, and a family of linearly degenerate cones $R(t) \subset \mathbb{P}^{n+2}$ over cubics such that the Schubert integral constructed for Case (C) in Section 3.4 is $Y_2 = \overline{\cup_{t \in \mathbb{C}} X_{w^2} \cap R(t)}$.*

In Cases (a-c), $[Y_1] = 2\xi_w$. In Case (d), $[Y_2] = 4\xi_w$.

Proof of Proposition 4.2(a.i). As noted in Section 2.5.1,

$$X_w = \mathbb{P}\langle e_1, \ldots, e_{\mathsf{j}} \rangle = \mathbb{P}^{\mathsf{j}-1} .$$

Following the notation of Section 3.3, set $C_1 = e^1_{\mathsf{j}} - e^{n+\mathsf{j}}_{n+1} \in \mathfrak{g}_{-\gamma}$ and $B = e^{\mathsf{j}}_{\mathsf{j}+1} - e^{n+\mathsf{j}+1}_{n+\mathsf{j}} \in \mathfrak{g}_{-\beta}$. Then $C_2 = [B, C_1] = e^1_{\mathsf{j}+1} - e^{n+\mathsf{j}+1}_{n+1}$ and $C_3 = [B, C_2] = 0$. Thus, the integer of (3.14) is $\mathsf{s} = 2$. By (3.16),

$$\mathfrak{m}_1 \;=\; \left\{ \textstyle\sum_{k=2}^{\mathsf{j}-1} s_k (e^1_k - e^{n+k}_{n+1}) \;+\; t(e^1_{\mathsf{j}} - e^{n+\mathsf{j}}_{n+1}) \;+\; \frac{1}{2} t^2 (e^1_{\mathsf{j}+1} - e^{n+\mathsf{j}+1}_{n+1}) \mid s_k, t \in \mathbb{C} \right\} .$$

With $o = [1 : 0 : \cdots : 0] \in \mathcal{Q}^m \subset \mathbb{P}^{m+1}$, we have

$$M_1 \cdot o = \left\{ [1 : s_2 : \cdots : s_{\mathsf{j}-1} : t : \tfrac{1}{2} t^2 : 0 : \cdots : 0] \in \mathbb{P}^{2n} \mid s_k, t \in \mathbb{C} \right\} .$$

The Schubert variety of Proposition 3.15 is

$$X_{w^1} = \mathcal{Q}^m \cap \mathbb{P}\langle e_1, \ldots, e_{\mathsf{j}+1} \rangle = \mathbb{P}^{\mathsf{j}} .$$

Let $Q \subset \mathbb{P}^{\mathsf{j}}$ be the quadric $\{0 = 2x_1 x_{\mathsf{j}+1} - (x_{\mathsf{j}})^2\}$. Then Lemma 3.17 yields $Y_1 = Q$. Thus $[Y_1] = 2\xi_w$. □

Proof of Proposition 4.2(a.ii). As above $X_w = \mathbb{P}\langle e_1, \ldots, e_n \rangle = \mathbb{P}^{n-1}$. Set $C_1 = e^1_n - e^{2n}_{n+1} \in \mathfrak{g}_{-\gamma}$ and $B = e^n_{2n+1} - e^{2n+1}_{2n} \in \mathfrak{g}_{-\beta}$. Then $C_2 = [B, C_1] = e^1_{2n+1} - e^{2n+1}_{n+1}$, $C_3 = [B, C_2] =$

[†] In [18, Section 6.1] we erroneously claimed that H_1 also fails when $\mathsf{j} = n$. As indicated by Case (c), it is H_2 that fails when $\mathsf{j} = n$.

$e^n_{n+1} - e^1_{2n}$ and $C_4 = [B, C_3] = 0$. Thus, the integer of (3.14) is $\mathsf{s} = 3$. By (3.16),

$$\begin{aligned} \mathfrak{m}_1 \; = \; \Big\{ \textstyle\sum_{k=2}^{n-1} s_k (e^1_k - e^{n+k}_{n+1}) \; &+ \; t(e^1_n - e^{2n}_{n+1}) \, + \, \tfrac{1}{2} t^2 (e^1_{2n+1} - e^{2n+1}_{n+1}) \\ &+ \; \tfrac{1}{6} t^3 (e^n_{n+1} - e^1_{2n}) \mid s_k, t \in \mathbb{C} \Big\} . \end{aligned}$$

So

$$M_1 \cdot o = \left\{ [1 : s_2 : \cdots : s_{n-1} : t : \tfrac{1}{24} t^4 : 0 : \cdots : 0 : -\tfrac{1}{6} t^3 : \tfrac{1}{2} t^2] \in \mathbb{P}^{2n} \mid s_k, t \in \mathbb{C} \right\} .$$

Set $\mathbb{P}^{n+2} = \mathbb{P}\langle e_1, \ldots, e_{n+1}, e_{2n}, e_{2n+1} \rangle$. The Schubert variety of Proposition 3.15 is $X_{w^1} = \mathcal{Q}^m \cap \mathbb{P}^{n+2}$. Let $Q = \{2x_1 x_{2n+1} - (x_n)^2 = 0\} \subset \mathbb{P}^{n+2}$ and $R = \{6(x_1)^2 x_{2n} + (x_n)^3 = 0\} \subset \mathbb{P}^{n+2}$. Then Lemma 3.17 yields $Y_1 = X_{w^1} \cap Q \cap R$. Intersecting Y_1 with a general Schubert variety representing the Poincaré dual of ξ_w, we find $[Y_1] = 2\xi_w$. ☐

Proof of Proposition 4.2(b). Define $\mathbb{P}^{2n-\mathtt{j}+1} = \mathbb{P}\langle e_1, \ldots, e_{n+1}, e_{n+\mathtt{j}+1}, \ldots, e_{2n+1} \rangle$. As noted in Section 2.5.1, $X_w = \mathcal{Q}^m \cap \mathbb{P}^{2n-\mathtt{j}+1}$.

Set $C_1 = e^1_{n+\mathtt{j}+1} - e^{\mathtt{j}+1}_{n+1}$ and $B = e^{n+\mathtt{j}+1}_{n+\mathtt{j}} - e^{\mathtt{j}}_{\mathtt{j}+1}$. Then $C_2 = [B, C_1] = e^1_{n+\mathtt{j}} - e^{\mathtt{j}}_{n+1}$ and $C_3 = [B, C_2] = 0$. So the integer of (3.14) is $\mathsf{s} = 2$, and (3.16) yields

$$\begin{aligned} \mathfrak{m}_1 \; = \; \Big\{ &\textstyle\sum_{k=2}^{n} s_k (e^1_k - e^{n+k}_{n+1}) \, + \, \sum_{\ell=\mathtt{j}+2}^{n} s'_\ell (e^1_{n+\ell} - e^\ell_{n+1}) \, + \, s'_{n+1} (e^1_{2n+1} - e^{2n+1}_{n+1}) \\ &+ t(e^1_{n+\mathtt{j}+1} - e^{\mathtt{j}+1}_{n+1}) \, + \, \tfrac{1}{2} t^2 (e^1_{n+\mathtt{j}} - e^{\mathtt{j}}_{n+1}) \mid s_k, s'_\ell, t \in \mathbb{C} \Big\} . \end{aligned}$$

So

$$\begin{aligned} M_1 \cdot o \;\; = \;\; \big\{ &[1 : s_2 : \cdots : s_n : \tau : 0 : \cdots 0 : \tfrac{1}{2} t^2 : t : s'_{\mathtt{j}+2} : \cdots : s'_{2n+1}] \in \mathbb{P}^{2n} \\ &\qquad\qquad\qquad\qquad \mid s_k, s'_\ell, t \in \mathbb{C} \big\} , \end{aligned}$$

with τ defined by $0 = 2\tau + (s'_{n+1})^2 + t^2 s_{\mathtt{j}} + 2t s_{\mathtt{j}+1} + 2 \sum_{k=\mathtt{j}+2}^{n} s_k s'_k$. Define $\mathbb{P}^{2n-\mathtt{j}+2} = \mathbb{P}\langle e_1, \ldots, e_{n+1}, e_{n+\mathtt{j}}, \ldots, e_{2n+1} \rangle$. The Schubert variety of Proposition 3.15 is $X_{w^1} = \mathcal{Q}^m \cap \mathbb{P}^{2n-\mathtt{j}+2}$. Let $Q = \{2x_1 x_{n+\mathtt{j}} - (x_{n+\mathtt{j}+1})^2 = 0\} \subset \mathbb{P}^{2n-\mathtt{j}+2}$. Lemma 3.17 yields $Y_1 = X_{w^1} \cap Q$. Intersecting Y_1 with a general Schubert variety representing the Poincaré dual of ξ_w, we find $[Y_1] = 2\xi_w$. ☐

Proof of Proposition 4.2(c). We have

$$\mathfrak{n}_w \; = \; \left\{ \textstyle\sum_{k=2}^{n} s_k (e^1_k - e^{n+k}_{n+1}) \, + \, t(e^1_{2n+1} - e^{2n+1}_{n+1}) \mid s_k, t \in \mathbb{C} \right\} .$$

Define $\mathbb{P}^{n+1} = \mathbb{P}\langle e_1, \ldots, e_{n+1}, e_{2n+1} \rangle$. Then $X_w = \overline{\exp(\mathfrak{n}_w) \cdot o} = \mathcal{Q}^m \cap \mathbb{P}^{n+1}$.

Following Section 3.5, set $C = e^1_{2n+1} - e^{2n+1}_{n+1} \in \mathfrak{g}_{-\gamma}$ and $E = e^n_1 - e^{n+1}_{2n} \in \mathfrak{g}_\varepsilon$. Then $B = [E, C] = e^{2n+1}_{2n} - e^n_{2n+1}$ and $C_\flat = [B, C] = e^1_{2n} - e^n_{n+1}$. Whence, the Schubert variety of Proposition 3.31 is $X_{w^2} = \mathcal{Q}^m \cap \mathbb{P}^{n+2}$, where $\mathbb{P}^{n+2} = \langle e_1, \ldots, e_{n+1}, e_{2n}, e_{2n+1} \rangle$. Additionally, $\mathrm{C}_2(t) = t(e^1_{2n+1} - e^{2n+1}_{n+1}) - \frac{1}{3} t^3 (e^1_{2n} - e^n_{n+1})$ and $\mathbf{b}(t) = \exp \frac{1}{2} t^2 (e^{2n+1}_{2n} - e^n_{2n+1})$. Thus

$$\exp\left(\mathrm{C}_2(t) + \mathfrak{n}_{w'}\right) \cdot (1, 0, \ldots, 0) \; = \; \left(1, s_2, \ldots, s_n, \tfrac{1}{3} t^3 s_n - \tfrac{1}{2} t^2, 0, \ldots, 0, -\tfrac{1}{3} t^3, t\right) \tag{4.3}$$

Fix $t \in \mathbb{C}$. Define $e_a(t) = \mathbf{b}(t)\cdot e_a$ for all $1 \le a \le 2n+1$. Then $e_a(t) = e_a$, if $a \neq n, 2n+1$, otherwise $e_n(t) = e_n - \frac{1}{2}t^2 e_{2n+1} - \frac{1}{8}t^4 e_{2n}$ and $e_{2n+1}(t) = e_{2n+1} + \frac{1}{2}t^2 e_{2n}$. Note that $\mathbf{b}(t)$ preserves $\mathbb{P}^{n+2} = \mathbb{P}\langle e_1, \dots, e_{n+1}, e_{2n}, e_{2n+1}\rangle$, and therefore preserves $X_{w^2} = \mathcal{Q}^m \cap \mathbb{P}^{n+2}$ (as asserted by Lemma 3.42). Let $[x_1 : \dots : x_{n+1} : x_{2n} : x_{2n+1}]$ denote the linear homogeneous coordinates on $\mathbb{P}^{n+2}$ associated with the basis $\{e_1(t), \dots, e_{n+1}(t), e_{2n}(t), e_{2n+1}(t)\}$, and define a linearly degenerate cubic $R(t) = \{0 = tx_1 - x_{2n+1}\} \cap \{0 = 3(x_1)^2 x_{2n} + (x_{2n+1})^3\} \subset \mathbb{P}^{n+2}$. From (3.38) and (4.3), we see that $Y_2(t) = \mathcal{Q}^m \cap R(t) = X_{w^2} \cap R(t)$. It now follows from Lemma 3.36 that $Y_2 = \overline{\cup_{t\in\mathbb{C}} Y_2(t)}$. One may check that $Y_2(\infty) := \lim_{|t|\to\infty} Y_2(t)$ is the line spanned by e_{2n}. Intersecting with a general Schubert variety representing the Poincaré dual of ξ_w yields $[Y_2] = 4\xi_w$. □

4.1.2. *Even dimensional quadrics* $\mathcal{Q}^{2n-2} = D_n/P_1$. Set $m = 2n-2$. Let ξ_w be a Schubert class on $\mathcal{Q}^m$, with associated pair $(\mathtt{a}, \mathtt{J})$; see Section 2.5.2. In [18, Section 6.2], we showed that H_1 is satisfied if and only if $\mathtt{a} = 0$ and $\mathtt{J} = \{\mathtt{j}\} \subset \{n-1, n\}$. In particular, H_1 fails when

(A) $\mathtt{a} = 0$ and $\mathtt{J} = \{\mathtt{j}\} \subset \{1, \dots, n-2\}$, for the pair $\gamma = \alpha_1 + \cdots + \alpha_{\mathtt{j}-1}$ and $\beta = \alpha_{\mathtt{j}}$;

(B) $\mathtt{a} = 0$ and $\mathtt{J} = \{n-1, n\}$, for the pair $\gamma = \alpha_1 + \cdots + \alpha_{n-2}$ and $\beta = \alpha_{n-1}$ (or $\beta = \alpha_n$);

(C) $\mathtt{a} = 1$ and $\mathtt{J} = \{\mathtt{j}\} \subset \{1, \dots, n-2\}$, for the pair $\gamma = \alpha_1 + \cdots + \alpha_{\mathtt{j}} + 2(\alpha_{\mathtt{j}+1} + \cdots + \alpha_{n-2}) + \alpha_{n-1} + \alpha_n$ and $\beta = \alpha_{\mathtt{j}}$;

(D) $\mathtt{a} = 1$ and $\mathtt{J} = \{n-1, n\}$, for the pairs $\gamma = \alpha_1 + \cdots + \alpha_{n-1}$ and $\beta = \alpha_n$, and $\gamma = \alpha_1 + \cdots + \alpha_{n-2} + \alpha_n$ and $\beta = \alpha_{n-1}$.

Lemma 4.4. *The condition* H_2 *holds for all* $w \in W^{\mathfrak{p}}$.

Lemma 4.4 is proved at the end of the section.

Proposition 4.5. *The nontrivial Schubert integrals constructed in Section 3.2 satisfy* $[Y_1] = 2\xi_w$. *Moreover,*

(i) *in Cases (A) and (B),* $X_w = \mathbb{P}^{\mathtt{j}-1}$ *and* $Y_1 = Q \subset \mathbb{P}^{\mathtt{j}} = X_{w^1}$ *is a cone over a conic with vertex* $\mathbb{P}^{\mathtt{j}-3}$*; and*

(ii) *in Cases (C) and (D),* $X_{w^1} = \mathcal{Q}^m \cap \mathbb{P}^{2n-\mathtt{j}+1}$ *contains* $X_w = \mathcal{Q}^m \cap \mathbb{P}^{2n-\mathtt{j}}$ *as a divisor and there exists a cone* $Q \subset \mathbb{P}^{2n-\mathtt{j}+1}$ *over a conic, with vertex* $\mathbb{P}^{2n-\mathtt{j}-2}$*, such that* $Y_1 = X_{w^1} \cap Q$.

Proof of Proposition 4.5(i). In Case (B) we may take either $\beta = \alpha_{n-1}$ or $\beta = \alpha_n$. The two cases are symmetric; without loss of generality we will prove the proposition for $\beta = \alpha_{n-1}$. Then Cases (A) and (B) may be uniformly expressed as: the condition H_1 fails for $\gamma = \alpha_1 + \cdots + \alpha_{\mathtt{j}-1}$ and $\beta = \alpha_{\mathtt{j}}$ with $2 \le \mathtt{j} \le n-1$. We have $\mathfrak{n}_w = \{\sum_{k=2}^{\mathtt{j}-1} s_k(e^1_k - e^{n+k}_{n+1}) + t(e^1_{\mathtt{j}} - e^{n+\mathtt{j}}_{n+1}) \mid s_k, t \in \mathbb{C}\}$. Set $\mathbb{P}^{\mathtt{j}-1} = \mathbb{P}\langle e_1, \dots, e_{\mathtt{j}}\rangle$. Then, (2.12) yields $X_w = \overline{\exp(\mathfrak{n}_w)\cdot o} = \mathcal{Q}^m \cap \mathbb{P}^{\mathtt{j}-1}$.

Following Section 3.3, we set $C_1 = e^1_{\mathtt{j}} - e^{n+\mathtt{j}}_{n+1} \in \mathfrak{g}_{-\gamma}$ and $B = e^{\mathtt{j}}_{\mathtt{j}+1} - e^{n+\mathtt{j}+1}_{n+\mathtt{j}} \in \mathfrak{g}_{-\beta}$. Then $C_2 = [B, C_1] = e^1_{\mathtt{j}+1} - e^{n+\mathtt{j}+1}_{n+1}$ and $C_3 = [B, C_2] = 0$. In particular, the integer of (3.14) is $\mathsf{s} = 2$. Equation (3.16) yields $\mathfrak{m}_1 = \{\sum_{k=2}^{\mathtt{j}-1} s_k(e^1_k - e^{n+k}_{n+1}) + t(e^1_{\mathtt{j}} - e^{n+\mathtt{j}}_{n+1}) + \frac{1}{2}t^2(e^1_{\mathtt{j}+1} - e^{n+\mathtt{j}+1}_{n+1}) \mid s_k, t \in \mathbb{C}\}$. Therefore,

$$M_1 \cdot o \;=\; \left\{[1 : s_2 : \cdots : s_{\mathtt{j}-1} : t : \tfrac{1}{2}t^2 : 0 : \cdots : 0] \mid s_k, t \in \mathbb{C}\right\} \;\subset\; \mathbb{P}^{2n-1}\,.$$

Set $\mathbb{P}^{\mathtt{j}} = \mathbb{P}\langle e_1, \dots, e_{\mathtt{j}+1}\rangle$. The Schubert variety of Proposition 3.15 is $X_{w^1} = \mathbb{P}^{\mathtt{j}}$. Let $Q = \{2x_1 x_{\mathtt{j}+1} - (x_{\mathtt{j}})^2 = 0\} \subset \mathbb{P}^{\mathtt{j}}$. Lemma 3.17 yields $Y_1 = Q$.

Intersecting Y_1 with a general Schubert variety representing the Poincaré dual of ξ_w yields $[Y_1] = 2\xi_w$. □

Proof of Proposition 4.5(ii). In Case (C) we may take either $\beta = \alpha_{n-1}$ or $\beta = \alpha_n$. The two cases are symmetric; without loss of generality we will prove the proposition for $\beta = \alpha_{n-1}$. To that end, set $2 \le \mathtt{j} \le n-2$ for Case (C), and $\mathtt{j} = n-1$ for Case (D). Then $\mathfrak{n}_w = \{\sum_{k=2}^n s_k(e^1_k - e^{n+k}_{n+1}) + \sum_{\ell=\mathtt{j}+2}^n s'_\ell(e^1_{n+\ell} - e^\ell_{n+1}) + t(e^1_{n+\mathtt{j}+1} - e^{\mathtt{j}+1}_{n+1}) \mid s_k, s'_\ell, t \in \mathbb{C}\}$, and (2.12) yields $X_w = \mathcal{Q}^m \cap \mathbb{P}^{2n-\mathtt{j}}$, where $\mathbb{P}^{2n-\mathtt{j}} = \mathbb{P}\langle e_1, \dots, e_{n+1}, e_{n+\mathtt{j}+1}, \dots, e_{2n}\rangle$.

Following Section 3.3, we set $C_1 = e^1_{n+\mathtt{j}+1} - e^{\mathtt{j}+1}_{n+1} \in \mathfrak{g}_{-\gamma}$ and $B = e^{n+\mathtt{j}+1}_{n+\mathtt{j}} - e^{\mathtt{j}}_{\mathtt{j}+1} \in \mathfrak{g}_{-\beta}$. Then $C_2 = [B, C_1] = e^1_{n+\mathtt{j}} - e^{\mathtt{j}}_{n+1}$ and $C_3 = [B, C_2] = 0$. Thus, the integer of (3.14) is $\mathsf{s} = 2$. By (3.16), we have $\mathfrak{m}_1 = \{\sum_{k=2}^n s_k(e^1_k - e^{n+k}_{n+1}) + \sum_{\ell=\mathtt{j}+2}^n s'_\ell(e^1_{n+\ell} - e^\ell_{n+1}) + t(e^1_{n+\mathtt{j}+1} - e^{\mathtt{j}+1}_{n+1}) + \frac{1}{2}t^2(e^1_{n+\mathtt{j}} - e^{\mathtt{j}}_{n+1}) \mid s_k, s'_\ell, t \in \mathbb{C}\}$. Whence

$$M_1 \cdot o \;=\; \left\{[1 : s_2 : \cdots : s_n : \tau : 0 : \cdots : 0 : \tfrac{1}{2}t^2 : t : s'_{\mathtt{j}+2} : \cdots : s'_n] \mid s_k, s'_\ell, t \in \mathbb{C}\right\} ,$$

where τ is given by $0 = 2\tau + t^2 s_{\mathtt{j}} + 2ts_{\mathtt{j}+1} + 2\sum_{\ell=\mathtt{j}+2}^n s_\ell s'_\ell$. Set $\mathbb{P}^{2n-\mathtt{j}+1} = \mathbb{P}\langle e_1, \dots, e_{n+1}, e_{n+\mathtt{j}}, \dots, e_{2n}\rangle$. The Schubert variety of Proposition 3.15 is $X_{w^1} = \mathcal{Q}^m \cap \mathbb{P}^{2n-\mathtt{j}+1}$. Define $Q = \{2x_1 x_{n+\mathtt{j}} - (x_{n+\mathtt{j}+1})^2 = 0\} \subset \mathbb{P}^{2n-\mathtt{j}+1}$. Then Lemma 3.17 yields $Y_1 = X_{w^1} \cap Q$.

Intersecting Y_1 with a general Schubert variety representing the Poincaré dual of ξ_w yields $[Y_1] = 2\xi_w$. □

Proof of Lemma 4.4. By Section 2.5.2, $\mathsf{a} \in \{0, 1\}$. By Remark 3.7, H_2 holds trivially when $\mathsf{a} = 0$. So suppose that $\mathsf{a} = 1$. By Section 2.5.2, there are two cases to consider $\mathtt{J} = \{\mathtt{j}\} \subset \{2, \dots, n-2\}$ and $\mathtt{J} = \{n-1, n\}$.

If $\mathtt{J} = \{\mathtt{j}\}$, then $\mathfrak{g}_{1,1}$ is an irreducible $\mathfrak{g}_{0,0}$–module with highest weight $\gamma = \alpha_1 + \cdots + \alpha_{\mathtt{j}} + 2(\alpha_{\mathtt{j}+1} + \cdots + \alpha_{n-2}) + \alpha_{n-1} + \alpha_n$. Similarly, $\mathfrak{g}_{1,0}$ is an irreducible $\mathfrak{g}_{0,0}$–module with highest weight $\varepsilon = \alpha_1 + \cdots + \alpha_{\mathtt{j}-1}$. Note that $\gamma' = \alpha_1 + \cdots + \alpha_n \in \Delta(\mathfrak{g}_{1,1})$. Since both $\gamma - \varepsilon$ and $\gamma' - \varepsilon$ are roots, (3.6.ii) fails. Thus, H_2 holds.

If $\mathtt{J} = \{n-1, n\}$, then $\mathfrak{g}_{1,1} = \mathfrak{g}_{\gamma_1} \oplus \mathfrak{g}_{\gamma_2}$ and both

$$\gamma_1 \;=\; \alpha_1 + \cdots + \alpha_{n-1} \quad \text{and} \quad \gamma_2 \;=\; \alpha_1 + \cdots + \alpha_{n-2} + \alpha_n$$

are highest $\mathfrak{g}_{0,0}$–weights. The $\mathfrak{g}_{0,0}$–module $\mathfrak{g}_{1,0}$ is irreducible with highest weight $\varepsilon = \alpha_1 + \cdots + \alpha_{n-2}$. Since both $\gamma_1 - \varepsilon = \alpha_{n-1}$ and $\gamma_2 - \varepsilon = \alpha_n$ are roots, (3.6.ii) fails. Thus, H_2 holds. □

4.2. Grassmannians.

We assume the notations and conventions of Sections 2.6 and 2.7 throughout this section. From Theorem 3.1 and [18, Theorem 6.1(a)], we obtain

Theorem 4.6. *Let $w \ne 1, w_0$ be an element of the Hasse diagram of $\mathrm{Gr}(\mathtt{i}, n+1)$ with corresponding data $(\mathsf{a}, \mathtt{J})$. Then the Schubert system $\mathcal{B}_w$ is rigid if and only if*

$$1 \;<\; \mathtt{j}_\ell - \mathtt{j}_{\ell+1} \quad \text{and} \quad 1 \;<\; \mathtt{k}_{m+1} - \mathtt{k}_m \tag{4.7}$$

holds for the following index ranges:

- ○ *if $\mathsf{p} = \mathsf{a}$, then $1 \le m \le \mathsf{a}$; and if $\mathsf{p} = \mathsf{a} + 1$, then $0 \le m \le \mathsf{a}$;*
- ○ *if $\mathsf{q} = \mathsf{a}$, then $1 \le \ell \le \mathsf{a}$; and if $\mathsf{q} = \mathsf{a} + 1$, then $0 \le \ell \le \mathsf{a}$.*

Remark (Partition formulation). Let $\lambda = \mu^{\mathsf{p}} \cdots \mu^0$ be the partition associated with w by Lemma 2.25. Then $\mathtt{j}_\ell - \mathtt{j}_{\ell+1}$ is the length of the block μ^ℓ, and $\mathtt{k}_{m+1} - \mathtt{k}_m$ is the length of the gap between the blocks μ^ℓ and $\mu^{\ell+1}$, when $\ell + m = \mathsf{a} + 1$.

Lemma 4.8. *The condition H_2 holds for all Schubert varieties in the Grassmannian.*

Proof. By Remark 3.7, the condition H_2 is satisfied if $\mathsf{a} = 0$. So suppose that $\mathsf{a} > 0$. Any highest $\mathfrak{g}_{0,0}$–weight $\varepsilon \in \Delta(\mathfrak{g}_{1,1-\mathsf{a}})$ is of the form

$$\varepsilon \ = \ \alpha_{\mathtt{j}_\ell+1} + \cdots + \alpha_{\mathtt{k}_m-1}$$

for $0 < \ell, m$ with $\ell + m = \mathsf{a} + 1$. Set

$$\begin{aligned} \gamma_1 &= \alpha_{\mathtt{j}_{\ell+1}+1} + \cdots + \alpha_{\mathtt{k}_m-1}\,, \\ \gamma_2 &= \alpha_{\mathtt{j}_\ell+1} + \cdots + \alpha_{\mathtt{k}_{m+1}-1}\,. \end{aligned}$$

Then $\gamma_1, \gamma_2 \in \Delta(\mathfrak{g}_{1,\mathsf{a}})$ are highest $\mathfrak{g}_{0,0}$–weights, and both $\gamma_1 - \varepsilon = \alpha_{\mathtt{j}_{\ell+1}+1} + \cdots + \alpha_{\mathtt{j}_\ell}$ and $\gamma_2 - \varepsilon = \alpha_{\mathtt{k}_m} + \cdots + \alpha_{\mathtt{k}_{m+1}-1}$ are roots. So (3.6.ii) fails, implying that H_2 holds. □

Suppose that the Schubert system $\mathcal{B}_w$ is flexible. By Theorem 4.6, either $1 = \mathtt{j}_\ell - \mathtt{j}_{\ell+1}$, for some ℓ, or $1 = \mathtt{k}_{m+1} - \mathtt{k}_m$, for some m. Let Y_1 be a nontrivial integral constructed in Section 3.2. In order to determine the integer $r > 0$ such that $[Y_1] = r\xi_w$, it suffices to consider the case that $1 = \mathtt{j}_\ell - \mathtt{j}_{\ell+1}$, for the case that $1 = \mathtt{k}_{m+1} - \mathtt{k}_m$ follows from the duality $\phi : \mathrm{Gr}(\mathtt{i}, n+1) \to \mathrm{Gr}(n+1-\mathtt{i}, n+1)$ mapping $E \subset \mathbb{C}^{n+1}$ to $\mathrm{Ann}(E) \subset \mathrm{Hom}(\mathbb{C}^{n+1}, \mathbb{C})$. To see this it suffices to observe that the map sends a Schubert variety X_w with description $\mathsf{a}(w) = \mathsf{a}$ and $\mathtt{J}(w) = \{\mathtt{j}_\mathtt{p}, \ldots, \mathtt{j}_1, \mathtt{k}_1, \ldots, \mathtt{k}_\mathtt{q}\}$ to the Schubert variety $\phi(X_w) = X_{w'}$ with description $\mathsf{a}(w') = \mathsf{a}$ and $\mathtt{J}(w') = \{\mathtt{j}'_\mathtt{q}, \ldots, \mathtt{j}'_1, \mathtt{k}'_1, \ldots, \mathtt{k}'_\mathtt{p}\}$, where $\mathtt{j}'_m = (n+1) - \mathtt{k}_m$ and $\mathtt{k}'_\ell = (n+1) - \mathtt{j}_\ell$. (See [18, Section 3.5].) Moreover, $Y_1 \subset \mathrm{Gr}(\mathtt{i}, n+1)$ is an integral of $\mathcal{B}_w$ if and only if $\phi(Y_1) \subset \mathrm{Gr}(n+1-\mathtt{i}, n+1)$ is an integral of $\mathcal{B}_{w'}$.

Proposition 4.9. *Fix $w \in W^{\mathfrak{p}} \backslash \{\mathbb{1}, w_0\}$. Let $\lambda = (\lambda_1, \ldots, \lambda_\mathtt{i})$ be the partition associated with w by Lemma 2.25, and let $F^\bullet$ be the full flag of (2.20). Suppose that $1 = \mathtt{j}_\ell - \mathtt{j}_{\ell+1}$ for some $\mathsf{a} + 1 - \mathsf{q} \le \ell \le \mathsf{a}$. Set $\mathsf{j} = \mathtt{j}_\ell$. The partition λ^1 associated with the Schubert variety X_{w^1} of Proposition 3.15 is given by $\lambda^1_\mathsf{j} = \lambda_\mathsf{j} + 1$, and $\lambda^1_s = \lambda_s$, for all $s \neq \mathsf{j}$. Moreover, there exists a cone $Q \subset F^{\lambda_\mathsf{j}+1}$ over a conic (with vertex $F^{\lambda_\mathsf{j}-2}$) such that the nontrivial integral constructed in Section 3.2 is*

$$Y_1 \ = \ \{E \in X_{w^1} \mid \exists\, \zeta \in \mathrm{Gr}(\mathsf{j}, E) \ \textit{s.t.}\ \zeta \subset Q\}\,. \tag{4.10}$$

In particular, $[Y_1] = 2\xi_w$.

Remark. I. Coskun [8] has observed that if Q is replaced with a cone over a curve of degree d, then the resulting variety (4.10) is irreducible and homologous to $d\,\xi_w$.

Proof. Set $\mathsf{k} = \mathtt{k}_{\mathsf{a}+1-\ell}$. The matrix representing $\mathfrak{n}_w$ is of the form

$$\mathfrak{n}_w \ = \ \begin{pmatrix} & & \vdots & \vdots & \\ \cdots & 0 & s^{\mathsf{j}}_{\mathsf{k}} & s^{\mathsf{j}+1}_{\mathsf{k}} & \cdots \\ \cdots & 0 & 0 & s^{\mathsf{j}+1}_{\mathsf{k}+1} & \cdots \\ \cdots & 0 & 0 & \vdots & \end{pmatrix}.$$

The condition H_1 fails for $\gamma = \alpha_{\mathsf{j}} + \cdots + \alpha_{\mathsf{k}-1}$ and $\beta = \alpha_{\mathsf{k}}$. The corresponding root spaces are spanned by $C := e^{\mathsf{j}}_{\mathsf{k}}$ and $B := -e^{\mathsf{k}}_{\mathsf{k}+1}$. We have $C_2 = e^{\mathsf{j}}_{\mathsf{k}+1}$ and $C_3 = 0$. In particular, the integer of (3.14) is $\mathsf{s} = 2$. Setting $t = s^{\mathsf{j}}_{\mathsf{k}}$, (3.16) yields

$$\mathfrak{m}_1 \;=\; \left(\begin{array}{ccccc} & & \vdots & \vdots & \\ \cdots & 0 & t & s^{\mathsf{j}+1}_{\mathsf{k}} & \cdots \\ \cdots & 0 & \tfrac{1}{2}t^2 & s^{\mathsf{j}+1}_{\mathsf{k}+1} & \cdots \\ \cdots & 0 & 0 & \vdots & \\ & \vdots & \vdots & \vdots & \end{array} \right).$$

Let Q be the rank three quadric in $\langle e_1, \ldots, e_{\mathsf{j}}, e_{\mathtt{i}+1}, \ldots, e_{\mathsf{k}+1} \rangle = \langle \mathsf{F}_\ell, e_{\mathsf{k}+1} \rangle = F^{\lambda_{\mathsf{j}}+1} = F^{\lambda^1_{\mathsf{j}}}$ given by $(x_{\mathsf{k}})^2 - 2x_{\mathsf{j}}x_{\mathsf{k}+1} = 0$. Proposition 3.15 and Lemma 3.17 yield (4.10).

By intersecting Y_1 with a general Schubert variety representing the Poincaré dual, we see that $[Y_1] = 2\xi_w$. □

4.3. Lagrangian grassmannians.

We assume the notations and conventions of Sections 2.8 and 2.9 throughout this section. Theorem 3.1 and [18, Theorem 6.1(c)] yield

Theorem 4.11. *Let $w \neq 1, w_0$ be an element of the Hasse diagram of $\mathrm{LG}(n, 2n)$ with corresponding data (a, J). Set $\mathsf{a}_0 = \mathsf{a} + 1 - \mathsf{p} \in \{0, 1\}$. The Schubert system $\mathcal{B}_w$ is rigid if and only if $1 < \mathsf{j}_\ell - \mathsf{j}_{\ell+1}$ for all $\mathsf{a}_0 \leq \ell \leq \mathsf{a}$.*

Remark (Partition formulation). Alternatively, if $\lambda = \mu^{\mathsf{p}} \cdots \mu^0$ is the partition associated with w (cf. Lemma 2.31), then the Schubert system is rigid if and only if the length $\mathsf{j}_\ell - \mathsf{j}_{\ell+1}$ of the block μ^ℓ is greater than 1 for all $\mathsf{a}_0 \leq \ell \leq \mathsf{a}$.

Lemma 4.12. *In the Lagrangian grassmannian the condition H_1 holds for a Schubert variety characterized (a, J) if and only if $1 < \mathsf{j}_\ell - \mathsf{j}_{\ell+1}$ for all $\mathsf{a}_0 \leq \ell \leq \mathsf{a}$ such that $\mathsf{a} \neq 2\ell - 1$. The condition H_2 holds if and only if $1 < \mathsf{j}_\ell - \mathsf{j}_{\ell+1}$ when $\mathsf{a} = 2\ell - 1$.*

The lemma is proved in [18, Section 6.3]. Proposition 4.13 describes the nontrivial integrals constructed in Sections 3.2 and 3.4.

Proposition 4.13. *Fix $w \in W^{\mathsf{p}} \backslash \{\mathbb{1}, w_0\}$. Let $\lambda = (\lambda_1, \ldots, \lambda_n)$ be the partition associated with w by Lemma 2.31, and let $F^\bullet$ be the full flag of (2.28).*

(a) *Suppose H_1 fails because $1 = \mathsf{j}_\ell - \mathsf{j}_{\ell+1}$ for some ℓ with $\mathsf{a}_0 \leq \ell \leq \mathsf{a}$ and $\mathsf{a} \neq 2\ell, 2\ell - 1$. Set $\mathsf{j} = \mathsf{j}_\ell$ and $\mathsf{k} = \mathsf{j}_{\mathsf{a}+1-\ell}$. The partition λ^1 associated with the Schubert variety X_{w^1} of Proposition 3.15 is given by $\lambda^1_i = \lambda_i + 1$, if $i = \mathsf{j}, \mathsf{k}$; otherwise $\lambda^1_i = \lambda_i$. There exists a cone $Q \subset F^{\lambda_{\mathsf{j}}+1}$ over a conic (with vertex $F^{\lambda_{\mathsf{j}}-2}$) such that the nontrivial integral constructed in Section 3.2 is*

$$Y_1 \;=\; \overline{\{E \in X_{w^1} \mid \exists \zeta \in \mathrm{Gr}(\mathsf{j}, E) \ s.t.\ \zeta \subset Q\}}\,. \tag{4.14a}$$

In particular, $[Y_1] = 2\xi_w$.

(b) *Suppose H_1 fails because $1 = \mathsf{j}_\ell - \mathsf{j}_{\ell+1}$ with $\mathsf{a} = 2\ell$ and $\mathsf{a}_0 \leq \ell \leq \mathsf{a}$. Set $\mathsf{j} = \mathsf{j}_\ell = \mathsf{j}_{\ell+1} + 1$. The partition λ^1 associated with the Schubert variety X_{w^1} of Proposition 3.15 is given by $\lambda^1_i = \lambda_i$, for all $i \neq \mathsf{j} - 1, \mathsf{j}$, and $\lambda^1_{\mathsf{j}} = \lambda_{\mathsf{j}} + 1 = n + 2$ and $\lambda^1_{\mathsf{j}-1} = \lambda_{\mathsf{j}-1} + 2 = n + 1$. There exist*

(i) *a cone* $Q \subset F^{n+2}$ *over a conic (with vertex* F^{n-1}*), and*

(ii) *a hyperplane* $H \subset F^{n+2}$ *(with* $\dim H \cap F^{n+1} = n$*), and a cone* $R \subset H$ *over a cubic curve (with vertex* F^{n-2}*)*

such that the nontrivial integral constructed in Section 3.2 is

$$Y_1 \;=\; \overline{\{E \in X_{w^1} \mid \exists\, \zeta_1, \zeta_2 \in \mathrm{Gr}(\mathsf{j}-1, E) \ s.t.\ \zeta_1 \subset Q\,,\ \zeta_2 \subset R\}}\,. \tag{4.14b}$$

Moreover, $[Y_1] = 2\xi_w$.

(c) *Suppose* H_2 *fails because* $1 = \mathsf{j}_\ell - \mathsf{j}_{\ell+1}$ *with* $\mathsf{a} = 2\ell - 1$*. Set* $\mathsf{j} := \mathsf{j}_\ell = \mathsf{j}_{\ell+1} + 1$*. The partition* λ^2 *associated with the Schubert variety* X_{w^2} *of Proposition 3.31 is given by* $\lambda^2_{\mathsf{j}} = \lambda_{\mathsf{j}} + 1 = n+1$*, otherwise* $\lambda^2_i = \lambda_i$*. There exist a curve* $\mathbf{b}(t)$ *in* $\mathrm{Stab}\, X_{w^2} \subset D_n$ *and linearly degenerate cones* $R(t) \subset \mathbf{b}(t) \cdot F^{n+1}$ *over cubic curves (with common vertices* F^{n-2}*) such that the nontrivial integral constructed in Section 3.4 is*

$$Y_2 \;=\; \overline{\bigcup_{t \in \mathbb{C}} Y_2(t)} \quad \text{with} \quad Y_2(t) \;=\; \overline{\{E \in X_{w^2} \mid \exists\, \zeta_t \in \mathrm{Gr}(\mathsf{j}, E) \ s.t.\ \zeta_t \subset R(t)\}}\,. \tag{4.14c}$$

Moreover, $[Y_2] = 3\xi_w$.

The proof is interspersed with examples illustrating the three cases of the proposition.

Example 4.15 (Illustration of Proposition 4.13(a)). Consider the Schubert variety $X_w \subset \mathrm{LG}(5, 10)$ corresponding to $\mathsf{a} = 3$ and $\mathsf{J} = \{1, 3, 4\}$. The Z_w–degrees of the root vectors in $\mathfrak{g}_{-1}$ are represented by the matrix

$$-\left[\begin{array}{c|cc|c|c} 6 & 5 & 5 & 4 & 3 \\ \hline 5 & 4 & 4 & 3 & 2 \\ 5 & 4 & 4 & 3 & 2 \\ \hline 4 & 3 & 3 & 2 & 1 \\ \hline 3 & 2 & 2 & 1 & 0 \end{array}\right].$$

So we identify $\mathfrak{n}_w \subset \mathfrak{g}_{-1}$ with the subspace

$$\mathfrak{n}_w \;=\; \begin{pmatrix} 0 & 0 & 0 & 0 & s^6_5 \\ 0 & 0 & 0 & s^7_4 & s^7_5 \\ 0 & 0 & 0 & s^8_4 & s^8_5 \\ 0 & s^9_2 & s^9_3 & s^9_4 & s^9_5 \\ s^{10}_1 & s^{10}_2 & s^{10}_3 & s^{10}_4 & s^{10}_5 \end{pmatrix},$$

with $s^{i+5}_j = s^{j+5}_i$. Set $F^2 = \langle e_1, e_{10}\rangle$, $F^5 = \langle F^2, e_2, e_3, e_9\rangle$ and $F^8 = \langle F^5, e_4, e_7, e_8\rangle$. Then

$$X_w \;=\; \{E \in \mathrm{LG}(5,10) \mid \dim(E \cap F^2) \geq 1\,,\ \dim(E \cap F^5) \geq 3\,,\ \dim(E \cap F^8) \geq 4\}\,.$$

The condition H_1 fails for $\gamma = \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5$ and $\beta = \alpha_1$. Set $C = C_1 = e^2_9 + e^4_7$ and $B = e^7_6 - e^1_2$. Then $C_2 = e^4_6 + e^1_9$ and $C_3 = 0$. In particular, the integer of (3.14) is $\mathsf{s} = 2$, and (3.16) yields

$$\mathfrak{m}_1 \;=\; \begin{pmatrix} 0 & 0 & 0 & \frac{1}{2}t^2 & s^6_5 \\ 0 & 0 & 0 & t & s^7_5 \\ 0 & 0 & 0 & s^8_4 & s^8_5 \\ \frac{1}{2}t^2 & t & s^9_3 & s^9_4 & s^9_5 \\ s^{10}_1 & s^{10}_2 & s^{10}_3 & s^{10}_4 & s^{10}_5 \end{pmatrix}.$$

The Schubert variety X_{w^1} of Proposition 3.15 is

$$X_{w^1} \;=\; \{E \in \mathrm{LG}(5,10) \mid \dim(E \cap F^5) \geq 3\}\,.$$

Let $Q \subset \langle F^8, e_6\rangle$ denote the rank three quadric defined by $\{(x_7)^2 - 2x_4x_6 = 0\}$. Lemma 3.17 yields

$$Y_1 \;=\; \overline{\{E \in X_{w^1} \mid \exists\, \zeta \in \mathrm{Gr}(4,E) \text{ s.t. } \zeta \subset Q\}}\,.$$

Intersecting Y_1 with a general Schubert variety representing the Poincaré dual ξ_w^* yields $[Y] = 2\xi_w$.

Proof of Proposition 4.13(a). There are two cases to consider.

Case 1: $\ell + 1 < m$. Set $\mathsf{k} := \mathsf{j}_m < \mathsf{j}_\ell =: \mathsf{j}$. Note that $\mathsf{k} < \mathsf{j}_{\ell+1} = \mathsf{j}_\ell - 1$. The condition H_1 fails for $\gamma = \alpha_{\mathsf{k}+1} + \cdots + \alpha_{\mathsf{j}-1} + 2(\alpha_{\mathsf{j}} + \cdots + n-1) + \alpha_n$ and $\beta = \alpha_{\mathsf{k}}$. The matrix representation of $\mathfrak{n}_w$ is

$$\mathfrak{n}_w \;=\; \begin{pmatrix}
 & & & & \vdots & \vdots & \vdots & \\
 & & & & 0 & 0 & s^{n+\mathsf{k}}_{\mathsf{j}+1} & \cdots \\
 & & & & 0 & s^{n+\mathsf{k}+1}_{\mathsf{j}} & s^{n+\mathsf{k}+1}_{\mathsf{j}+1} & \cdots \\
 & & & & & \vdots & \vdots & \\
\cdots & 0 & 0 & & & \vdots & \vdots & \\
\cdots & 0 & s^{n+\mathsf{j}}_{\mathsf{k}+1} & \cdots & \cdots & s^{n+\mathsf{j}}_{\mathsf{j}} & s^{n+\mathsf{j}}_{\mathsf{j}+1} & \cdots \\
\cdots & s^{n+\mathsf{j}+1}_{\mathsf{k}} & s^{n+\mathsf{j}+1}_{\mathsf{k}+1} & \cdots & \cdots & s^{n+\mathsf{j}+1}_{\mathsf{j}} & s^{n+\mathsf{j}+1}_{\mathsf{j}+1} & \cdots \\
 & \vdots & \vdots & & & \vdots & \vdots &
\end{pmatrix}.$$

Set $C_1 = e^{\mathsf{k}+1}_{n+\mathsf{j}} + e^{\mathsf{j}}_{n+\mathsf{k}+1}$ and $B = e^{n+\mathsf{k}+1}_{n+\mathsf{k}} - e^{\mathsf{k}}_{\mathsf{k}+1}$. Then $C_2 = [B, C_1] = e^{\mathsf{j}}_{n+\mathsf{k}} + e^{\mathsf{k}}_{n+\mathsf{j}}$ and $C_3 = 0$. Whence, the integer of (3.14) is $\mathsf{s} = 2$. Setting $t = s^{n+\mathsf{j}}_{\mathsf{k}+1}$, (3.16) yields

$$\mathfrak{m}_1 \;=\; \begin{pmatrix}
 & & & & 0 & 0 & \vdots & \\
 & & & & 0 & \frac{1}{2}t^2 & s^{n+\mathsf{k}}_{\mathsf{j}+1} & \cdots \\
 & & & & 0 & t & s^{n+\mathsf{k}+1}_{\mathsf{j}+1} & \cdots \\
 & & & & & \vdots & \vdots & \\
0 & 0 & 0 & & & \vdots & \vdots & \\
0 & \frac{1}{2}t^2 & t & \cdots & \cdots & s^{n+\mathsf{j}}_{\mathsf{j}} & s^{n+\mathsf{j}}_{\mathsf{j}+1} & \cdots \\
\cdots & s^{n+\mathsf{j}+1}_{\mathsf{k}} & s^{n+\mathsf{j}+1}_{\mathsf{k}+1} & \cdots & \cdots & s^{n+\mathsf{j}+1}_{\mathsf{j}} & s^{n+\mathsf{j}+1}_{\mathsf{j}+1} & \cdots \\
 & \vdots & \vdots & & & \vdots & \vdots &
\end{pmatrix}.$$

By (2.26a),

$$F^{\lambda_{\mathsf{j}}+1} \;=\; \langle \mathtt{F}_\ell, e_{n+\mathsf{k}}\rangle \;=\; \langle e_1, \ldots, e_{\mathsf{j}}\,,\, e_{n+\mathsf{k}}, \ldots, e_{2n}\rangle\,.$$

Let $Q = \{2x_{\mathsf{j}}x_{n+\mathsf{k}} - (x_{n+\mathsf{k}+1})^2 = 0\} \subset F^{\lambda_{\mathsf{j}}+1}$. It follows from Proposition 3.15 and Lemma 3.17 that Y_1 is given by (4.14a).

Intersecting Y_1 with a general Schubert variety representing the Poincaré dual ξ_w^* yields $[Y] = 2\xi_w$.

Case 2: $m < \ell$. Set $\mathsf{j} := \mathsf{j}_\ell < \mathsf{j}_m =: \mathsf{k}$. Then H_1 fails for $\gamma = \alpha_{\mathsf{j}} + \cdots + \alpha_{\mathsf{k}} + 2(\alpha_{\mathsf{k}+1} + \cdots + \alpha_{n-1}) + \alpha_n$ and $\beta = \alpha_{\mathsf{k}}$. In this case

$$\mathfrak{n}_w \;=\; \left(\begin{array}{cccccccc}
 & & & & & \vdots & \vdots & \\
 & & & & & 0 & 0 & \\
 & & & & & 0 & s^{n+\mathsf{j}}_{\mathsf{k}+1} & \cdots \\
 & & & & & s^{n+\mathsf{j}+1}_{\mathsf{k}} & s^{n+\mathsf{j}+1}_{\mathsf{k}+1} & \cdots \\
 & & & & & \vdots & \vdots & \\
\cdots & 0 & 0 & s^{n+\mathsf{k}}_{\mathsf{j}+1} & \cdots & s^{n+\mathsf{k}}_{\mathsf{k}} & s^{n+\mathsf{k}}_{\mathsf{k}+1} & \cdots \\
\cdots & 0 & s^{n+\mathsf{k}+1}_{\mathsf{j}} & s^{n+\mathsf{k}+1}_{\mathsf{j}+1} & \cdots & s^{n+\mathsf{k}+1}_{\mathsf{k}} & s^{n+\mathsf{k}+1}_{\mathsf{k}+1} & \cdots \\
 & & \vdots & \vdots & & \vdots & \vdots &
\end{array}\right).$$

Set $C = C_1 = e^{\mathsf{j}}_{n+\mathsf{k}+1} + e^{\mathsf{k}+1}_{n+\mathsf{j}}$ and $B = e^{n+\mathsf{k}+1}_{n+\mathsf{k}} - e^{\mathsf{k}}_{\mathsf{k}+1}$. Then $C_2 = [B, C_1] = e^{\mathsf{j}}_{n+\mathsf{k}} + e^{\mathsf{k}}_{n+\mathsf{j}}$ and $C_3 = 0$. In particular, the integer of (3.14) is $\mathsf{s} = 2$ and (3.16) yields

$$\mathfrak{m}_1 \;=\; \left(\begin{array}{cccccccc}
 & & & & & \vdots & \vdots & \\
 & & & & & 0 & 0 & \\
 & & & & 0 & \frac{1}{2}t^2 & t & \cdots \\
 & & & & \cdots & s^{n+\mathsf{j}+1}_{\mathsf{k}} & s^{n+\mathsf{j}+1}_{\mathsf{k}+1} & \cdots \\
 & & 0 & \vdots & & \vdots & \vdots & \\
\cdots & 0 & \frac{1}{2}t^2 & s^{n+\mathsf{k}}_{\mathsf{j}+1} & \cdots & s^{n+\mathsf{k}}_{\mathsf{k}} & s^{n+\mathsf{k}}_{\mathsf{k}+1} & \cdots \\
\cdots & 0 & t & s^{n+\mathsf{k}+1}_{\mathsf{j}+1} & \cdots & s^{n+\mathsf{k}+1}_{\mathsf{k}} & s^{n+\mathsf{k}+1}_{\mathsf{k}+1} & \cdots \\
 & & \vdots & \vdots & & \vdots & \vdots &
\end{array}\right).$$

Recall the definition (2.26a) of $\mathtt{F}_\ell$, and let $Q \subset \langle \mathtt{F}_\ell, e_{n+\mathsf{k}} \rangle = F^{\lambda_{\mathsf{j}}+1}$ be the quadric defined by $2x_{\mathsf{j}} x_{n+\mathsf{k}} - (x_{n+\mathsf{k}+1})^2 = 0$. It follows from Proposition 3.15 and Lemma 3.17 that the integral Y_1 is given by (4.14a).

Intersecting Y_1 with a general Schubert variety representing the Poincaré dual ξ_w^* yields $[Y_1] = 2\xi_w$. □

Example 4.16 (Illustration of Proposition 4.13(b)). Consider the Schubert variety $X_w \subset \mathrm{LG}(4,8)$ corresponding to $\mathtt{a} = 2$ and $\mathtt{J} = \{2,3\}$. The Z_w–degrees of the root vectors in $\mathfrak{g}_{-1}$ are represented by the matrix

$$-\left[\begin{array}{cc|c|c}
4 & 4 & 3 & 2 \\
4 & 4 & 3 & 2 \\
\hline
3 & 3 & 2 & 1 \\
\hline
2 & 2 & 1 & 0
\end{array}\right].$$

So we identify $\mathfrak{n}_w \subset \mathfrak{g}_{-1}$ with the subspace

$$\mathfrak{n}_w \ = \ \begin{pmatrix} 0 & 0 & 0 & s^5_4 \\ 0 & 0 & 0 & s^6_4 \\ 0 & 0 & s^7_3 & s^7_4 \\ s^8_1 & s^8_2 & s^8_3 & s^8_4 \end{pmatrix} ,$$

with $s^{i+4}_j = s^{j+4}_i$. Set $F^3 = \langle e_1, e_2, e_8 \rangle$ and $F^5 = \langle F^3, e_3, e_7 \rangle$. Then

$$X_w \ = \ \{ E \in \mathrm{LG}(4,8) \mid \dim(E \cap F^3) \geq 2 \, , \ \dim(E \cap F^5) \geq 3 \} \, .$$

The condition H_1 fails for $\gamma = 2\alpha_3 + \alpha_4$ and $\beta = \alpha_2$. Set $C_1 = e^3_7$ and $B = e^6_7 - e^3_2$. Then $C_2 = [B, C_1] = e^3_6 + e^2_7$, $C_3 = e^2_6$ and $C_4 = 0$. Thus, the integer of (3.14) is $\mathsf{s} = 3$, and (3.16) yields

$$\mathfrak{m}_1 \ = \ \begin{pmatrix} 0 & 0 & 0 & s^5_4 \\ 0 & \frac{1}{6}t^3 & \frac{1}{2}t^2 & s^6_4 \\ 0 & \frac{1}{2}t^2 & t & s^7_4 \\ s^8_1 & s^8_2 & s^8_3 & s^8_4 \end{pmatrix} ,$$

Set $F^2 = \langle e_1, e_8 \rangle$, $F^5 = \langle F^3, e_6, e_7 \rangle$ and $F^6 = \langle F^5, e_3 \rangle$. The Schubert variety of Proposition 3.15 is

$$X_{w^1} \ = \ \{ E \in \mathrm{LG}(4,8) \mid \dim(E \cap F^2) \geq 1 \, , \ \dim(E \cap F^6) \geq 3 \} \, .$$

Let $Q \subset F^6$ be the rank three quadric $\{ (x_7)^2 - 2x_3x_6 = 0 \}$, and let $R \subset F^5$ be the cubic $\{ x_2(6x_6)^2 - (2x_7)^3 = 0 \}$. Lemma 3.17 yields

$$Y_1 \ = \ \overline{\{ E \in X_{w^1} \mid \exists \, \zeta_1, \zeta_2 \in \mathrm{Gr}(2,E) \text{ s.t. } \zeta_1 \subset Q \, , \ \zeta_2 \subset R \}} \, .$$

Intersecting Y_1 with a general Schubert variety representing the Poincaré dual ξ^*_w yields $[Y] = 2\xi_w$.

Proof of Proposition 4.13(b). Here we have

$$\mathfrak{n}_w \ = \ \begin{pmatrix} & & & \vdots & \vdots & \\ \cdots & 0 & 0 & 0 & s^{n+\mathsf{j}-1}_{\mathsf{j}+1} & \cdots \\ \cdots & 0 & 0 & s^{n+\mathsf{j}}_{\mathsf{j}} & s^{n+\mathsf{j}}_{\mathsf{j}+1} & \cdots \\ \cdots & \cdots & s^{n+\mathsf{j}+1}_{\mathsf{j}-1} & s^{n+\mathsf{j}+1}_{\mathsf{j}} & s^{n+\mathsf{j}+1}_{\mathsf{j}+1} & \cdots \\ & & \vdots & \vdots & \vdots & \ddots \end{pmatrix} .$$

The condition H_1 fails for $\gamma = 2(\alpha_{\mathsf{j}} + \cdots + \alpha_{n-1}) + \alpha_n$ and $\beta = \alpha_{\mathsf{j}-1}$. Set $C = C_1 = e^{\mathsf{j}}_{n+\mathsf{j}}$ and $B = e^{n+\mathsf{j}}_{n+\mathsf{j}+1} - e^{\mathsf{j}-1}_{\mathsf{j}}$. Then $C_2 = [B, C_1] = e^{\mathsf{j}}_{n+\mathsf{j}-1} + e^{\mathsf{j}-1}_{n+\mathsf{j}}$, $C_3 = [B, C_2] = 2e^{\mathsf{j}-1}_{n+\mathsf{j}-1}$ and $C_4 = 0$. Thus, the integer of (3.14) is $\mathsf{s} = 3$, and (3.16) yields

$$\mathfrak{m}_1 \ = \ \begin{pmatrix} & 0 & 0 & 0 & \vdots & \\ \cdots & 0 & \frac{1}{3}t^3 & \frac{1}{2}t^2 & s^{n+\mathsf{j}-1}_{\mathsf{j}+1} & \cdots \\ \cdots & 0 & \frac{1}{2}t^2 & t & s^{n+\mathsf{j}}_{\mathsf{j}+1} & \cdots \\ \cdots & \cdots & s^{n+\mathsf{j}+1}_{\mathsf{j}-1} & s^{n+\mathsf{j}+1}_{\mathsf{j}} & s^{n+\mathsf{j}+1}_{\mathsf{j}+1} & \cdots \\ & & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

With $\mathtt{F}_\bullet$ and $F^\bullet$ as defined in (2.26a) and (2.28), we have

$$F^{n+2} \;=\; \langle \mathtt{F}_\ell, e_{n+\mathsf{j}-1}\rangle \;=\; \langle e_1, \ldots e_{\mathsf{j}}\,,\, e_{n+\mathsf{j}-1}, \ldots, e_{2n}\rangle\,.$$

Define

$$H \;=\; \langle \mathtt{F}_{\ell+1}, e_{n+\mathsf{j}-1}, e_{n+\mathsf{j}}\rangle \;=\; \langle e_1, \ldots, e_{\mathsf{j}-1}\,,\, e_{n+\mathsf{j}-1}, \ldots, e_{2n}\rangle \;=\; \langle F^n, e_{n+\mathsf{j}-1}\rangle\,.$$

Let $Q \subset F^{n+2}$ be the quadric hypersurface $2x_{\mathsf{j}}x_{n+\mathsf{j}-1} - (x_{n+\mathsf{j}})^2 = 0$, and let $R \subset H$ be the cubic hypersurface $x_{\mathsf{j}-1}(3x_{n+\mathsf{j}-1})^2 - (2x_{n+\mathsf{j}})^3 = 0$. Then Proposition 3.15 and Lemma 3.17 yield (4.14b).

Intersecting Y_1 with a general Schubert variety representing the Poincaré dual ξ_w^* yields $[Y_1] = 2\xi_w$. □

Example 4.17 (Illustration of Proposition 4.13(c)). Consider the Schubert variety $X_w \subset \mathrm{LG}(5,10)$ corresponding to $\mathsf{a} = 3$ and $\mathtt{J} = \{1,2,4\}$. The Z_w–degrees of the root vectors in $\mathfrak{g}_{-1}$ are represented by the matrix

$$-\left[\begin{array}{c|c|cc|c}
6 & 5 & 4 & 4 & 3 \\ \hline
5 & 4 & 3 & 3 & 2 \\ \hline
4 & 3 & 2 & 2 & 1 \\
4 & 3 & 2 & 2 & 1 \\ \hline
3 & 2 & 1 & 1 & 0
\end{array}\right].$$

So we identify $\mathfrak{n}_w \subset \mathfrak{g}_{-1}$ with the subspace

$$\mathfrak{n}_w \;=\; \begin{pmatrix}
0 & 0 & 0 & 0 & s^6_5 \\
0 & 0 & s^7_3 & s^7_4 & s^7_5 \\
0 & s^8_2 & s^8_3 & s^8_4 & s^8_5 \\
0 & s^9_2 & s^9_3 & s^9_4 & s^9_5 \\
s^{10}_1 & s^{10}_2 & s^{10}_3 & s^{10}_4 & s^{10}_5
\end{pmatrix},$$

with $s^{i+5}_j = s^{j+5}_i$. Set $F^2 = \langle e_1, e_{10}\rangle$, $F^5 = \langle F^2, e_2, e_8, e_9\rangle$ and $F^8 = \langle F^5, e_3, e_4, e_7\rangle$. Then

$$X_w \;=\; \{E \in \mathrm{LG}(5,10) \mid \dim(E\cap F^2) \geq 1\,,\ \dim(E\cap F^5) \geq 2\,,\ \dim(E \cap F^8) \geq 4\}\,.$$

The condition H_2 fails for $\gamma = \alpha_2 + 2(\alpha_3+\alpha_4) + \alpha_5$ and $\varepsilon = 2(\alpha_3+\alpha_4)+\alpha_5$. Set $C = e^3_7 + e^2_8$ and $E = -e^8_3$. Then $B = [E,C] = e^8_7 - e^2_3$ and $C_\flat = [B,C] = 2e^2_7$. (Here $\beta = \alpha_2$.) We have $\mathrm{C}_2(t) = t(e^3_7 + e^2_8) - \frac{2}{3}t^3 e^2_7$ and $\mathbf{b}(t) = \exp \frac{1}{2}t^2(e^8_7 - e^2_3)$. Fix $t \in \mathbb{C}$, and define $e_j(t) := \mathbf{b}(t)e_j$. Then $\{e_j(t)\}_{j=1}^{10}$ is a basis of $\mathbb{C}^{10}$. Explicitly, $e_j(t) = e_j$, for all $j \neq 2,8$; while $e_2(t) = e_2 - \frac{1}{2}t^2 e_3$ and $e_8(t) = e_8 + \frac{1}{2}t^2 e_7$. With respect to this basis, the subset $\mathfrak{m}_2(t) \subset \mathfrak{m}_2$ of (3.37) is expressed as

$$\mathfrak{m}_2(t) \;=\; \begin{pmatrix}
0 & 0 & 0 & 0 & s^6_5 \\
0 & -\frac{2}{3}t^3 & t & s^7_4 & s^7_5 \\
0 & t & s^8_3 & s^8_4 & s^8_5 \\
0 & s^9_2 & s^9_3 & s^9_4 & s^9_5 \\
s^{10}_1 & s^{10}_2 & s^{10}_3 & s^{10}_4 & s^{10}_5
\end{pmatrix}.$$

By Lemma 3.42, the Schubert variety of Proposition 3.31 is

$$X_{w^2} \;=\; \{E \in \mathrm{LG}(5,10) \mid \dim(E \cap F^2) \geq 1\,,\ \dim(E \cap F^8) \geq 4\}\,.$$

Set $F^6(t) = \mathbf{b}(t) \cdot \langle e_1, e_2, e_7, e_8, e_9, e_{10} \rangle = \langle e_1, e_2(t), e_7, e_8, e_9, e_{10} \rangle$, and let $[x_1 : x_2 : x_7 : x_8 : x_9 : x_{10}]$ be the linear homogeneous coordinates on $F^6(t)$ associated with the basis $\{e_j(t)\}$. Let $R(t) \subset F^6(t)$ be the linearly degenerate cubic defined by $\{0 = tx_2 - x_8 = 2(x_8)^3 + 3(x_2)^2 x_7\}$. Then (3.38) yields

$$Y_2(t) \ = \ \overline{\{E \in X_{w^2} \mid \exists \, \zeta_t \in \mathrm{Gr}(2, E) \text{ s.t. } \zeta_t \subset R(t)\}} \,.$$

By Proposition 3.31, $Y_2 = \overline{\cup_{t \in \mathbb{C}} Y_2(t)}$.

Remark. From (3.38), and the expression for $\mathfrak{m}_2(t)$ above, we see that $Y_2(\infty) := \lim_{|t| \to \infty} Y_2(t)$ is the set of all planes $E \in \mathrm{LG}(n, 2n)$ such that $\langle e_7 \rangle \subset E \subset \langle e_1, e_3, \ldots, e_{10} \rangle$, $\dim(E \cap \langle e_1, e_7, e_{10} \rangle) \geq 2$, $\dim(E \cap \langle e_1, e_3, e_7, \ldots, e_{10} \rangle) \geq 3$ and $\dim(E \cap \langle e_1, e_3, e_4, e_7, \ldots, e_{10} \rangle) \geq 4$.

Intersecting Y_2 with a general Schubert variety representing the Poincaré dual ξ_w^* yields $[Y_2] = 3\xi_w$.

Proof of Proposition 4.13(c). In this case, H_2 fails for $\gamma = \alpha_{\mathrm{j}} + 2(\alpha_{\mathrm{j}+1} + \cdots + \alpha_{n-1}) + \alpha_n$ and $\varepsilon = 2(\alpha_{\mathrm{j}+1} + \cdots + \alpha_{n-1}) + \alpha_n$.

$$\mathfrak{n}_w \ = \ \begin{pmatrix} & \vdots & \vdots & \vdots & \vdots & \\ \cdots & 0 & 0 & 0 & s^{n+\mathrm{j}-1}_{\mathrm{j}+2} & \cdots \\ \cdots & 0 & 0 & s^{n+\mathrm{j}}_{\mathrm{j}+1} & s^{n+\mathrm{j}}_{\mathrm{j}+2} & \cdots \\ \cdots & 0 & s^{n+\mathrm{j}+1}_{\mathrm{j}} & s^{n+\mathrm{j}+1}_{\mathrm{j}+1} & s^{n+\mathrm{j}+1}_{\mathrm{j}+2} & \cdots \\ \cdots & s^{n+\mathrm{j}+2}_{\mathrm{j}-1} & s^{n+\mathrm{j}+2}_{\mathrm{j}} & s^{n+\mathrm{j}+2}_{\mathrm{j}+1} & s^{n+\mathrm{j}+2}_{\mathrm{j}+2} & \cdots \\ & \vdots & \vdots & \vdots & \vdots & \end{pmatrix} .$$

Set $C = e^{\mathrm{j}}_{n+\mathrm{j}+1} + e^{\mathrm{j}+1}_{n+\mathrm{j}}$ and $E = -e^{n+\mathrm{j}+1}_{\mathrm{j}+1}$. Then $B = [E, C] = e^{n+\mathrm{j}+1}_{n+\mathrm{j}} - e^{\mathrm{j}}_{\mathrm{j}+1}$ and $C_\flat = [B, C] = 2e^{\mathrm{j}}_{n+\mathrm{j}}$. From (3.34), we have $\mathrm{C}_2(t) = t(e^{\mathrm{j}}_{n+\mathrm{j}+1} + e^{\mathrm{j}+1}_{n+\mathrm{j}}) - \frac{2}{3}t^3 e^{\mathrm{j}}_{n+\mathrm{j}}$, and $\mathbf{b}(t) = \exp \frac{1}{2} t^2 B$. Define $e_a(t) := \mathbf{b}(t) \cdot e_a$. With respect to the basis $\{e_a(t)\}_{a=1}^{2n}$ of $\mathbb{C}^{2n}$, the set $\mathfrak{m}_2(t)$ of (3.35) is represented by the matrix

$$\mathfrak{m}_2(t) \ = \ \begin{pmatrix} & \vdots & \vdots & \vdots & \vdots & \\ \cdots & 0 & 0 & 0 & s^{n+\mathrm{j}-1}_{\mathrm{j}+2} & \cdots \\ \cdots & 0 & -\frac{2}{3}t^3 & t & s^{n+\mathrm{j}}_{\mathrm{j}+2} & \cdots \\ \cdots & 0 & t & s^{n+\mathrm{j}+1}_{\mathrm{j}+1} & s^{n+\mathrm{j}+1}_{\mathrm{j}+2} & \cdots \\ \cdots & s^{n+\mathrm{j}+2}_{\mathrm{j}-1} & s^{n+\mathrm{j}+2}_{\mathrm{j}} & s^{n+\mathrm{j}+2}_{\mathrm{j}+1} & s^{n+\mathrm{j}+2}_{\mathrm{j}+2} & \cdots \\ & \vdots & \vdots & \vdots & \vdots & \end{pmatrix} .$$

By Lemma 3.42, $\mathbf{b}(t)$ stabilizes X_{w^2}. Consequently, the Schubert variety of Proposition 3.31 is

$$X_{w^2} \ = \ \{E \in \mathrm{LG}(n, 2n) \mid \dim(E \cap \mathtt{F}_i) \geq \mathtt{j}_i \ \forall \ i \neq \ell\} \,.$$

Fix $t \in \mathbb{C}$. With $F^\bullet$ as defined by (2.28), let $F^\bullet(t) = \mathbf{b}(t) \cdot F^\bullet$. Then

$$\begin{aligned} F^{n-2}(t) \ &= \ F^{n-2} \ = \ \langle e_1, \ldots, e_{\mathrm{j}-1} \,, e_{n+\mathrm{j}+2}, \ldots, e_{2n} \rangle \,, \\ F^{n+2}(t) \ &= \ F^{n+2} \ = \ \langle e_1, \ldots, e_{\mathrm{j}+1} \,, e_{n+\mathrm{j}}, \ldots, e_{2n} \rangle \end{aligned}$$

are constant. We have

$$\begin{aligned} F^n(t) &= \langle e_1, \ldots, e_{\mathsf{j}-1}\,,\, e_{\mathsf{j}} - \tfrac{1}{2}t^2 e_{\mathsf{j}+1}\,,\, e_{n+\mathsf{j}+1} + \tfrac{1}{2}t^2 e_{n+\mathsf{j}}\,,\, e_{n+\mathsf{j}+2}, \ldots, e_{2n}\rangle\,, \\ F^{n+1}(t) &= \langle e_1, \ldots, e_{\mathsf{j}-1}\,,\, e_{\mathsf{j}} - \tfrac{1}{2}t^2 e_{\mathsf{j}+1}\,,\, e_{n+\mathsf{j}}\,,\, e_{n+\mathsf{j}+1}\,,\, e_{n+\mathsf{j}+2}, \ldots, e_{2n}\rangle\,. \end{aligned}$$

Let x denote the linear homogeneous coordinates on $F^{n+1}(t)$ associated with the basis given by the $\{e_j(t)\}$. Let $R(t) \subset F^{n+1}(t)$ be the linearly degenerate cubic $\{x_{n+\mathsf{j}+1} = tx_{\mathsf{j}}\} \cap \{0 = 2(x_{n+\mathsf{j}+1})^3 + 3x_{n+\mathsf{j}}(x_{\mathsf{j}})^2\}$. Then (3.38) yields the expression for $Y_2(t)$ in (4.14c). From Proposition 3.31 we obtain the expression for Y_2 in (4.14c).

Intersecting Y_2 with a general Schubert variety representing the Poincaré dual ξ_w^* yields $[Y_2] = 3\xi_w$. □

4.4. **Spinor varieties.** The notations and conventions of Sections 2.10 and 2.11 hold throughout this section. By Theorem 3.1 and [18, Theorem 6.1], we have

Theorem 4.18. *Let* $w \in W^{\mathfrak{p}}\backslash\{\mathbb{1}, w_0\}$ *be an element of the Hasse diagram of* $\mathcal{S}_n$ *with corresponding data* (a, J)*. The Schubert system* $\mathcal{B}_w$ *is rigid if and only if* $|\mathsf{J}| \le \mathsf{a} + 1$*,* $1 < \mathsf{j}_\ell - \mathsf{j}_{\ell+1}$ *for all* $1 \le \ell \le \mathsf{a} + \alpha_{n-1}(Z_w)$*, and* $2 < \mathsf{j}_{\mathsf{r}} - \mathsf{j}_{\mathsf{r}+1}$ *when* $\mathsf{r} > \alpha_{n-1}(Z_w)$*.*

In [18, Section 6.4], we saw that there are three cases in which the condition H_1 may fail:

(A) $1 = \mathsf{j}_\ell - \mathsf{j}_{\ell+1}$ for some $1 \le \ell \le \mathsf{a} + \alpha_{n-1}(Z_w)$, with $\ell \ne \mathsf{r}$ when $\mathsf{r} > \alpha_{n-1}(Z_w)$;
(B) $2 = \mathsf{j}_{\mathsf{r}} - \mathsf{j}_{\mathsf{r}+1}$, when $\mathsf{r} > \alpha_{n-1}(Z_w)$; or
(C) $\mathsf{p} = \mathsf{a} + 2$.

Lemma 4.19. *The condition* H_2 *holds for all Schubert varieties in the Spinor variety* $\mathcal{S}_n = D_n/P_n$*.*

The lemma is proved at the end of the section. Definition 3.4 and Lemma 4.19 implies that H_+ fails if and only if H_1 fails. Equivalently, one of Case (a), (b) or (c) is satisfied.

Proposition 4.20. *Suppose that* $\mathcal{B}_w$ *is flexible,* $w \in W^{\mathfrak{p}}\backslash\{\mathbb{1}, w_0\}$*. Let* λ *and* λ^1 *respectively be the partitions associated with* X_w *and* X_{w^1} *(cf. Proposition 3.15) by Lemma 2.39, and let* $F^\bullet$ *be the full flag of Section 2.11. With the exceptions indicated below,* $\lambda_i^1 = \lambda_i$*.*

(a) *Suppose* $1 = \mathsf{j}_\ell - \mathsf{j}_{\ell+1}$ *for some* $1 \le \ell \le \mathsf{a} + \alpha_{n-1}(Z_w)$*, with* $\ell \ne \mathsf{r}$ *when* $\mathsf{r} > \alpha_{n-1}(Z_w)$*. Set* $\mathsf{j} = \mathsf{j}_\ell$ *and* $\mathsf{k} = \mathsf{j}_m$*, with* $\ell + m = \mathsf{a} + 1 + \alpha_{n-1}(Z_w)$*. Then* $\lambda_i^1 = \lambda_i + 1$*, if* $i = \mathsf{j}, \mathsf{k}$*.*
(b) *Suppose* $2 = \mathsf{j}_{\mathsf{r}} - \mathsf{j}_{\mathsf{r}+1}$ *and* $\mathsf{r} > \alpha_{n-1}(Z_w)$*.*
  (i) *If* $\mathsf{a} + \alpha_{n-1}(Z_w) = 2\mathsf{r} - 1$*, set* $\mathsf{j} = \mathsf{j}_{\mathsf{r}} - 1$ *and* $\mathsf{k} = \mathsf{j}_{\mathsf{r}}$*. Then* $\lambda_{\mathsf{j}}^1 = \lambda_{\mathsf{j}} + 1$ *and* $\lambda_{\mathsf{k}}^1 = \lambda_{\mathsf{k}} + 2$*.*
  (ii) *If* $\mathsf{a} + \alpha_{n-1}(Z_w) = 2\mathsf{r}$*, set* $\mathsf{j} = \mathsf{j}_{\mathsf{r}}$ *and* $\mathsf{k} = \mathsf{j}_{\mathsf{r}+1}$*. Then* $\lambda_i^1 = \lambda_i^1 + 1$ *if* $i = \mathsf{j}, \mathsf{k}$*.*
(c) *Suppose* $\mathsf{p} = \mathsf{a} + 2$*. Set* $\mathsf{j} = n$ *and* $\mathsf{k} = \mathsf{j}_{\mathsf{p}}$*. Then* $\lambda_i^1 = \lambda_i + 1$ *if* $i = \mathsf{j}, \mathsf{k}$*.*

There exists a cone $Q \subset F^{\lambda_{\mathsf{j}}+1}$ *over a conic (with vertex* $F^{\lambda_{\mathsf{j}}-2}$*) such that the nontrivial integral constructed in Section 3.2 is*

$$Y_1 = \overline{\{E \in X_{w^1} \mid \exists \zeta \in \mathrm{Gr}(\mathsf{j}, E) \ s.t.\ \zeta \subset Q\}}\,. \tag{4.21}$$

In particular, $[Y_1] = 2\xi_w$*.*

Remark (Partition formulation). Recall (Section 2.11) that the length of the block μ^ℓ is $\mathsf{j}_\ell - \mathsf{j}_{\ell+1}$. Also, in Case (b.i), $\mu^{\mathsf{r}} = (n-1, n)$; and in Case (b.ii), $\mu^{\mathsf{r}} = (n+1, n+2)$.

The proof of Proposition 4.20 is very similar to the proofs of Propositions 4.9 and 4.13, so I will sketch it with less detail than given in Sections 4.2 and 4.3.

Proof of Proposition 4.20 in Case (a). The argument breaks into two subcases: $\ell < \mathsf{r}$ and $\mathsf{r} < \ell$. In the first case, H_1 fails for

$$\gamma \ = \ \alpha_{\mathsf{k}+1} + \cdots + \alpha_{\mathsf{j}-1} + 2(\alpha_{\mathsf{j}} + \cdots + \alpha_{n-2}) + \alpha_{n-1} + \alpha_n \quad \text{and} \quad \beta \ = \ \alpha_{\mathsf{k}}\,.$$

In the second case, H_1 fails for

$$\gamma \ = \ \alpha_{\mathsf{j}} + \cdots + \alpha_{\mathsf{k}} + 2(\alpha_{\mathsf{k}+1} + \cdots + \alpha_{n-2}) + \alpha_{n-1} + \alpha_n \quad \text{and} \quad \beta \ = \ \alpha_{\mathsf{k}}\,.$$

In both cases we have $C_1 = e^{\mathsf{j}}_{n+\mathsf{k}+1} - e^{\mathsf{k}+1}_{n+\mathsf{j}}$ and $B = e^{n+\mathsf{k}+1}_{n+\mathsf{k}} - e^{\mathsf{k}}_{\mathsf{k}+1}$. Thus $C_2 = [B, C_1] = e^{\mathsf{j}}_{n+\mathsf{k}} - e^{\mathsf{k}}_{n+\mathsf{j}}$ and $C_3 = 0$. In particular, the integer of (3.14) is $\mathsf{s} = 2$. Let $Q \subset \langle e_1, \ldots, e_{\mathsf{j}}\,,\, e_{n+\mathsf{k}}, \ldots, e_{2n}\rangle = F^{\lambda_{\mathsf{j}}+1}$ be the quadric $\{0 = 2x_{\mathsf{j}}x_{n+\mathsf{k}} - (x_{n+\mathsf{k}+1})^2\}$. Then Proposition 3.15 and Lemma 3.17 imply that the nontrivial integral constructed in Section 3.2 is (4.21).

Intersecting Y_1 with a general Schubert variety representing the Poincaré dual of ξ_w yields $[Y_1] = 2\xi_w$. □

Proof of Proposition 4.20 in Case (b). The argument breaks into two subcases: $\mathsf{a}+\alpha_{n-1}(Z_w) = 2\mathsf{r}$ and $\mathsf{a} + \alpha_{n-1}(Z_w) = 2\mathsf{r} - 1$.

Subcase (i): $\mathsf{a} + \alpha_{n-1}(Z_w) = 2\mathsf{r} - 1$. (Note that $\mathsf{k} = \mathsf{j} + 1$.) The condition H_1 fails for the pair

$$\gamma \ = \ \alpha_{\mathsf{k}-1} + \alpha_{\mathsf{k}} + 2(\alpha_{\mathsf{k}+1} + \cdots + \alpha_{n-2}) + \alpha_{n-1} + \alpha_n \quad \text{and} \quad \beta \ = \ \alpha_{\mathsf{k}}\,.$$

Set $C_1 = e^{\mathsf{j}}_{n+\mathsf{k}+1} - e^{\mathsf{k}+1}_{n+\mathsf{j}}$ and $B = e^{n+\mathsf{k}+1}_{n+\mathsf{k}} - e^{\mathsf{k}}_{\mathsf{k}+1}$. Then $C_2 = [B, C_1] = e^{\mathsf{j}}_{n+\mathsf{k}} - e^{\mathsf{k}}_{n+\mathsf{j}}$ and $C_3 = 0$. In particular, the integer of (3.14) is $\mathsf{s} = 2$. Let $Q \subset \langle e_1, \ldots, e_{\mathsf{j}}\,,\, e_{n+\mathsf{k}}, \ldots, e_{2n}\rangle = F^{\lambda_{\mathsf{j}}+1}$ be the quadric $\{0 = 2x_{\mathsf{j}}x_{n+\mathsf{k}} - (x_{n+\mathsf{k}+1})^2\}$. Then Proposition 3.15 and Lemma 3.17 imply that the nontrivial integral constructed in Section 3.2 is (4.21).

Intersecting Y_1 with a general Schubert variety representing the Poincaré dual of ξ_w yields $[Y_1] = 2\xi_w$.

Subcase (ii): $\mathsf{a} + \alpha_{n-1}(Z_w) = 2\mathsf{r}$. (Note that $\mathsf{j} = \mathsf{k} + 2$.) The condition H_1 fails for the pair

$$\gamma \ = \ \alpha_{\mathsf{j}-1} + 2(\alpha_{\mathsf{j}} + \cdots + \alpha_{n-2}) + \alpha_{n-1} + \alpha_n \quad \text{and} \quad \beta \ = \ \alpha_{\mathsf{k}}\,.$$

Set $C_1 = e^{\mathsf{j}-1}_{n+\mathsf{j}} - e^{\mathsf{j}}_{n+\mathsf{j}-1}$ and $B = e^{n+\mathsf{k}+1}_{n+\mathsf{k}} - e^{\mathsf{k}}_{\mathsf{k}+1}$. Then $C_2 = [B, C_1] = e^{\mathsf{k}}_{n+\mathsf{j}} - e^{\mathsf{j}}_{n+\mathsf{k}}$ and $C_3 = 0$. In particular, the integer of (3.14) is $\mathsf{s} = 2$. Let $Q \subset \langle e_1, \ldots, e_{\mathsf{j}}\,,\, e_{n+\mathsf{k}}, \ldots, e_{2n}\rangle = F^{\lambda_{\mathsf{j}}+1}$ be the quadric $\{0 = 2x_{\mathsf{j}+1}x_{n+\mathsf{j}-1} - (x_{n+\mathsf{j}})^2\}$. Then Proposition 3.15 and Lemma 3.17 imply that the nontrivial integral constructed in Section 3.2 is (4.21).

Intersecting Y_1 with a general Schubert variety representing the Poincaré dual of ξ_w yields $[Y_1] = 2\xi_w$. □

Proof of Proposition 4.20 in Case (c). The condition H_1 fails for the pair

$$\gamma \ = \ \alpha_{\mathsf{k}+1} + \cdots + \alpha_{n-2} + \alpha_n \quad \text{and} \quad \beta \ = \ \alpha_{\mathsf{k}}\,.$$

Set $C_1 = e^n_{n+\mathsf{k}+1} - e^{\mathsf{k}+1}_{2n}$ and $B = e^{n+\mathsf{k}+1}_{n+\mathsf{k}} - e^{\mathsf{k}}_{\mathsf{k}+1}$. Then $C_2 = e^n_{n+\mathsf{k}} - e^{\mathsf{k}}_{2n}$ and $C_3 = 0$. In particular, the integer of (3.14) is $\mathsf{s} = 2$. Let $Q \subset \langle \mathtt{F}_0, e_{n+\mathsf{k}} \rangle = F^{\lambda_n + 1}$ be the quadric $\{0 = 2x_n x_{n+\mathsf{k}} - (x_{n+\mathsf{k}+1})^2\}$. Then Proposition 3.15 and Lemma 3.17 imply that the nontrivial integral constructed in Section 3.2 is (4.21).

Intersecting Y_1 with a general Schubert variety representing the Poincaré dual of ξ_w yields $[Y_1] = 2\xi_w$. $\square$

Proof of Lemma 4.19. Let a and J be the integer and marking characterizing a Schubert variety in $\mathcal{S}_n$. If $\mathsf{a} = 0$, then $\mathfrak{g}_{1,\mathsf{a}-1} = \{0\}$ and H_2 holds trivially. So suppose that $\mathsf{a} > 0$. If $n - 1 \notin \mathsf{J}$, then a highest $\mathfrak{g}_{0,0}$–weight $\varepsilon \in \Delta(\mathfrak{g}_{1,\mathsf{a}-1})$ is of the form

$$\varepsilon \ = \ \alpha_{\mathsf{j}_\ell + 1} + \cdots + \alpha_{\mathsf{j}_m} + 2(\alpha_{\mathsf{j}_m+1} + \cdots + \alpha_{n-2}) + \alpha_{n-1} + \alpha_n \,, \tag{4.22}$$

with $\ell + m = \mathsf{a} + 1$. If $n - 1 \in \mathsf{J}$, then a highest $\mathfrak{g}_{0,0}$–weight $\varepsilon \in \Delta(\mathfrak{g}_{1,\mathsf{a}-1})$ is either of the form (4.22), with $\ell + m = \mathsf{a} + 2$, or

$$\varepsilon \ = \ \alpha_{\mathsf{j}_\mathsf{a}+1} + \cdots + \alpha_{n-2} + \alpha_n \,. \tag{4.23}$$

If ε is of the form (4.22), then both

$$\begin{aligned} \gamma &= \alpha_{\mathsf{j}_{\ell+1}+1} + \cdots + \alpha_{\mathsf{j}_m} + 2(\alpha_{\mathsf{j}_m+1} + \cdots + \alpha_{n-2}) + \alpha_{n-1} + \alpha_n \,, \\ \gamma &= \alpha_{\mathsf{j}_\ell+1} + \cdots + \alpha_{\mathsf{j}_{m+1}} + 2(\alpha_{\mathsf{j}_{m+1}+1} + \cdots + \alpha_{n-2}) + \alpha_{n-1} + \alpha_n \,, \end{aligned}$$

are highest $\mathfrak{g}_{0,0}$–weights in $\Delta(\mathfrak{g}_{1,\mathsf{a}})$ with the property that $\gamma - \varepsilon \in \Delta$. In particular, (3.6.ii) cannot hold. If ε is of the form (4.23), then both

$$\gamma \ = \ \alpha_{\mathsf{j}_{\mathsf{a}+1}+1} + \cdots + \alpha_{n-2} + \alpha_n \quad \text{and} \quad \gamma \ = \ \alpha_{\mathsf{j}_\mathsf{a}+1} + \cdots + \alpha_n$$

are highest $\mathfrak{g}_{0,0}$–weights in $\Delta(\mathfrak{g}_{1,\mathsf{a}})$ with the property that $\gamma - \varepsilon \in \Delta$. Again, (3.6.ii) cannot hold. $\square$

4.5. **The exceptional cases.**

Theorem 4.24. *Suppose that X is either the Cayley plane E_6/P_6 or the Freudenthal variety E_7/P_7. The Schubert rigid classes ξ_w are precisely those associated to the nodes in Figures 1 and 2 that are circled twice.*

Proof. By [18, Theorem 6.1(e)], the twice circled nodes correspond to the Schubert varieties satisfying H_+. By Theorem 3.1, the Schubert system is rigid if and only if H_+ is satisfied. $\square$

Remark 4.25. I used [16] to verify that:

(a) The condition H_2 holds for all $w \in W^\mathfrak{p} \backslash \{\mathbb{1}, w_0\}$.

(b) If H_1 fails for $w \in W^\mathfrak{p} \backslash \{\mathbb{1}, w_0\}$, then the integer of (3.14) is $\mathsf{s} = 2$.

5. The Schur system

Assume that X is irreducible. (See [18, Table 1].) If X is classical, then the $w \in W^\mathfrak{p}$ for which $B_w \neq R_w$ are identified in [18, Section 8]. (Similar computations yield the corresponding result in the exceptional cases; see Figures 1 and 2.) However the results were not clearly stated. That is unfortunate as it is interesting to know when the Schur and Schubert systems coincide. (It is also an interesting question from the perspective of representation theory to determine when R_w consists of a single orbit B_w, see Remark

2.11(b).) In this section I will rectify this omission (see Theorems 5.1, 5.2 and 5.3, and Section 5.5).

5.1. **Grassmannian $\mathbf{Gr}(\mathtt{i}, \boldsymbol{n}+\mathbf{1}) = \boldsymbol{A_n}/\boldsymbol{P}_{\mathtt{i}}$.** Assume the notation and conventions of Section 2.6.

Theorem 5.1 ([18])**.** *Fix* $w \in W^{\mathfrak{p}} \backslash \{\mathbb{1}, w_0\}$ *with associated data* $(\mathsf{a}, \mathtt{J})$*. Then* $R_w \neq B_w$ *if and only if there exist* $\mathtt{j}_\ell, \mathtt{k}_m \in \mathtt{J}$ *satisfying one of the following:*

(a) $\ell + m = \mathsf{a} + 1$ *and* $1 \;=\; \mathtt{j}_\ell - \mathtt{j}_{\ell+1} \;=\; \mathtt{k}_{m+1} - \mathtt{k}_m$*; or*

(b) $\ell + m = \mathsf{a} + 2$ *and* $1 \;=\; \mathtt{j}_{\ell-1} - \mathtt{j}_\ell \;=\; \mathtt{k}_m - \mathtt{k}_{m-1}$.

Theorem 5.1 is proved in [18, Section 8.2].

Coskun has studied the rigidity and smoothability of Schubert classes on the Grassmannian. Theorems 1.3 and 1.6 of [6] are as follows.

Theorem (Coskun)**.**

(1) *The Schubert class* ξ_w *is flexible if and only if there exist* $\mathtt{j}_\ell, \mathtt{k}_m \in \mathtt{J}$ *such that the conditions of Theorem 5.1(a) hold.*

(2) *If there exists an index* $1 \leq \ell \leq \mathsf{a}$ *such that either* $1 < \mathtt{j}_\ell - \mathtt{j}_{\ell+1}$*, or* $1 < \mathtt{k}_{\ell+1} - \mathtt{k}_\ell$*, then the Schubert class* ξ_w *does not admit a smooth algebraic representative.*

5.2. Quadric hypersurfaces $\boldsymbol{\mathcal{Q}^m} \subset \mathbb{P}^{\boldsymbol{m}+\mathbf{1}}$.

5.2.1. *Odd dimensional quadrics* $\mathcal{Q}^{2n-1} = B_n/P_1$. Each $\bigwedge^\ell \mathfrak{g}_{-1}$ is an irreducible $\mathfrak{g}_0 = \mathbb{C} \oplus \mathfrak{b}_{n-1}$–module. So $R_w = \mathrm{Gr}(|w|, \mathfrak{g}_{-1})$. However, $B_w = G_0 \cdot \mathfrak{n}_w \subsetneqq \mathrm{Gr}(|w|, \mathfrak{g}_{-1})$. Thus $B_w \neq R_w$.

5.2.2. *Even dimensional quadrics* $\mathcal{Q}^{2n-2} = D_n/P_1$. If $|w| = n-1$, then $B_w = R_w$; see [18, Section 8.3].

Suppose that $|w| \neq n-1$. If $n > 4$, then, each $\bigwedge^\ell \mathfrak{g}_{-1}$ is an irreducible $\mathfrak{g}_0 = \mathbb{C} \oplus \mathfrak{d}_{n-1}$–module. However, $B_w = G_0 \cdot \mathfrak{n}_w \subsetneqq \mathrm{Gr}(|w|, \mathfrak{g}_{-1})$. Thus $B_w \neq R_w$.

When $n = 4$, the semi-simple part of the reductive $\mathfrak{g}_0$ is $\mathfrak{a}_3 = \mathfrak{sl}_4\mathbb{C}$ and $\mathfrak{g}_{-1} = \bigwedge^2 \mathbb{C}^4$. So G_0 acts transitively on $\mathrm{Gr}(1, \mathfrak{g}_{-1})$ implying $B_w = R_w$ for $(\mathsf{a}(w), \mathtt{J}(w)) = (0, \{2\})$, which corresponds to $X_w = \mathbb{P}^1$.

5.3. **Lagrangian Grassmannians $\mathbf{LG}(\boldsymbol{n}, \mathbf{2}\boldsymbol{n}) = \boldsymbol{C_n}/\boldsymbol{P_n}$.** Assume the notations and conventions of Section 2.8.

Theorem 5.2. *Fix* $w \in W^{\mathfrak{p}} \backslash \{\mathbb{1}, w_0\}$ *with associated data* $(\mathsf{a}, \mathtt{J})$*. Then* $R_w \neq B_w$ *if and only if there exists* $\mathtt{j}_\ell, \mathtt{j}_m \in \mathtt{J}$ *satisfying one of the following:*

(a) $\ell + m = \mathsf{a} + 1$ *and* $1 = \mathtt{j}_\ell - \mathtt{j}_{\ell+1} = \mathtt{j}_m - \mathtt{j}_{m+1}$*,*

(b) $\ell + m = \mathsf{a} + 2$ *and* $1 = \mathtt{j}_{\ell-1} - \mathtt{j}_\ell = \mathtt{j}_{m-1} - \mathtt{j}_m$.

The proof of Theorem 5.2 is given in [18, Section 8.4].

5.4. Spinor varieties $\mathcal{S}_n = D_n/P_n$.

Assume the notations and conventions of Section 2.10.

Theorem 5.3. *Fix $w \in W^{\mathfrak{p}} \backslash \{\mathbb{1}, w_0\}$ with associated data $(\mathtt{a}, \mathtt{J})$. Then $R_w \neq B_w$ if and only if there exists $\mathtt{j}_\ell, \mathtt{j}_m \in \mathtt{J}$ satisfying one of the following:*

(a) $\ell + m = \mathtt{a} + 1 + \alpha_{n-1}(Z_w)$ *and* $1 + \delta_{\ell m} = \mathtt{j}_\ell - \mathtt{j}_{\ell+1} = \mathtt{j}_m - \mathtt{j}_{m+1}$, *or*

(b) $\ell + m = \mathtt{a} + 2 + \alpha_{n-1}(Z_w)$ *and* $1 + \delta_{\ell m} = \mathtt{j}_{\ell-1} - \mathtt{j}_\ell = \mathtt{j}_{m-1} - \mathtt{j}_m$.

Above, $\delta_{\ell m}$ is the Kronecker delta. The proof of Theorem 5.3 is given in [18, Section 8.4].

5.5. The exceptional cases.

With the assistance of [16], the $w \in W^{\mathfrak{p}}$ satisfying $R_w = B_w$ may also be identified. See Figures 1 and 2.

References

[1] Sara Billey and V. Lakshmibai. *Singular loci of Schubert varieties*, volume 182 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 2000.

[2] Armand Borel. Kählerian coset spaces of semisimple Lie groups. *Proc. Nat. Acad. Sci. U. S. A.*, 40:1147–1151, 1954.

[3] Armand Borel and André Haefliger. La classe d'homologie fondamentale d'un espace analytique. *Bull. Soc. Math. France*, 89:461–513, 1961.

[4] R. L. Bryant. *Rigidity and Quasi-Rigidity of Extremal Cycles in Hermitian Symmetric Spaces*, volume 153 of *Annals of Mathematics Studies*. Princeton University Press, 2010. arXiv:math/0006186.

[5] Andreas Čap and Jan Slovák. *Parabolic geometries. I*, volume 154 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2009. Background and general theory.

[6] Izzet Coskun. Rigid and non-smoothable Schubert classes. *J. Differential Geom.*, 87(3):493–514, 2011.

[7] Izzet Coskun. Rigidity of schubert classes in orthogonal grassmannians, 2011. In preparation.

[8] Izzet Coskun, 2012. personal communication.

[9] Robin Hartshorne, Elmer Rees, and Emery Thomas. Nonsmoothing of algebraic cycles on Grassmann varieties. *Bull. Amer. Math. Soc.*, 80:847–851, 1974.

[10] Jaehyun Hong. Rigidity of singular Schubert varieties in $\mathrm{Gr}(m,n)$. *J. Differential Geom.*, 71(1):1–22, 2005.

[11] Jaehyun Hong. Rigidity of smooth Schubert varieties in Hermitian symmetric spaces. *Trans. Amer. Math. Soc.*, 359(5):2361–2381 (electronic), 2007.

[12] Steven L. Kleiman. The transversality of a general translate. *Compositio Math.*, 28:287–297, 1974.

[13] Anthony W. Knapp. *Lie groups beyond an introduction*, volume 140 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, second edition, 2002.

[14] Bertram Kostant. Lie algebra cohomology and the generalized Borel-Weil theorem. *Ann. of Math. (2)*, 74:329–387, 1961.

[15] Bertram Kostant. Lie algebra cohomology and generalized Schubert cells. *Ann. of Math. (2)*, 77:72–144, 1963.

[16] LiE. Computer algebra package for semisimple Lie algebra computations, www-math.univ-poitiers.fr/∼maavl/LiE/.

[17] C. Robles. Singular loci of cominuscule Schubert varieties. Preprint, 2012.

[18] C. Robles and D. The. Rigid schubert varieties in compact hermitian symmetric spaces. *Selecta Mathematica, New Series*, pages 1–61, 2012. 10.1007/s00029-011-0082-y.

[19] Dennis The, 2011. personal communication.

[20] Hugh Thomas and Alexander Yong. A combinatorial rule for (co)minuscule Schubert calculus. *Adv. Math.*, 222(2):596–620, 2009.

[21] Maria Walters. *Geometry and uniqueness of some extreme subvarieties in complex Grassmannians*. PhD thesis, University of Michigan, 1997.

E-mail address: `robles@math.tamu.edu`

Mathematics Department, Mail-stop 3368, Texas A&M University, College Station, TX 77843-3368